

ON WITT VECTOR COHOMOLOGY FOR SINGULAR VARIETIES

PIERRE BERTHELOT, SPENCER BLOCH, AND HÉLÈNE ESNAULT

ABSTRACT. Over a perfect field k of characteristic $p > 0$, we construct a “Witt vector cohomology with compact supports” for separated k -schemes of finite type, extending (after tensorisation with \mathbb{Q}) the classical theory for proper k -schemes. We define a canonical morphism from rigid cohomology with compact supports to Witt vector cohomology with compact supports, and we prove that it provides an identification between the latter and the slope < 1 part of the former. Over a finite field, this allows one to compute congruences for the number of rational points in special examples. In particular, the congruence modulo the cardinality of the finite field of the number of rational points of a theta divisor on an abelian variety does not depend on the choice of the theta divisor. This answers positively a question by J.-P. Serre.

CONTENTS

1. Introduction	1
2. Witt vector cohomology with compact supports	6
3. A descent theorem	17
4. Witt vector cohomology and rigid cohomology	21
5. Proof of the main theorem	28
6. Applications and examples	33
References	41

1. INTRODUCTION

Let k be a perfect field of characteristic $p > 0$, $W = W(k)$, $K = \text{Frac}(W)$. If X is a proper and smooth variety defined over k , the theory of the de Rham-Witt complex and the degeneration of the slope spectral sequence provide a functorial isomorphism ([6, III, 3.5], [20, II, 3.5])

$$(1.1) \quad H_{\text{crys}}^*(X/K)^{<1} \xrightarrow{\sim} H^*(X, W\mathcal{O}_X)_K,$$

where $H_{\text{crys}}^*(X/K)^{<1}$ is the maximal subspace on which Frobenius acts with slopes < 1 , and the subscript K denotes tensorisation with K .

If we only assume that X is proper, but maybe singular, we can use rigid cohomology to generalize crystalline cohomology while retaining all the standard properties of a topological cohomology theory. Thus, the left hand side of (1.1) remains well defined, and, when k is finite, the alternated product of the corresponding characteristic polynomials of Frobenius can be interpreted as the factor of the zeta function $\zeta(X, t)$ “of slopes < 1 ” (see 6.1 for a precise definition). On the other hand, the classical theory of the de Rham-Witt complex can no longer be directly applied to X , but the sheaf of Witt vectors $W\mathcal{O}_X$ is still available. Thus the right hand side of (1.1) remains also well defined. As in the smooth case, this is a finitely generated K -vector space endowed with a Frobenius action with slopes in $[0, 1[$ (Proposition 2.10), which has the advantage of being directly related to the coherent cohomology of X . It is therefore of interest to know whether, when X is singular, (1.1) can be generalized as an isomorphism

$$(1.2) \quad H_{\text{rig}}^*(X/K)^{<1} \xrightarrow{\sim} H^*(X, W\mathcal{O}_X)_K,$$

where $H_{\text{rig}}^*(X/K)^{<1}$ denotes the subspace of slope < 1 of $H_{\text{rig}}^*(X/K)$.

Our main result gives a positive answer to this question. More generally, we show that a “Witt vector cohomology with compact supports” can be defined for separated k -schemes of finite type, giving cohomology spaces $H_c^*(X, W\mathcal{O}_{X,K})$ which are finite dimensional K -vector spaces, endowed with a Frobenius action with slopes in $[0, 1[$. Then, for any such scheme X , the slope < 1 subspace of the rigid cohomology with compact supports of X has the following description:

Theorem 1.1. *Let k be a perfect field of characteristic $p > 0$, X a separated k -scheme of finite type. There exists a functorial isomorphism*

$$(1.3) \quad H_{\text{rig,c}}^*(X/K)^{<1} \xrightarrow{\sim} H_c^*(X, W\mathcal{O}_{X,K}).$$

This is a striking confirmation of Serre’s intuition [24] about the relation between topological and Witt vector cohomologies. On the other hand, this result bears some analogy with Hodge theory. Recall from [8] (or [12]) that if X is a proper scheme defined over \mathbb{C} , then its Betti cohomology $H^*(X, \mathbb{C})$ is a direct summand of its de Rham cohomology $\mathbb{H}^*(X, \Omega_X^\bullet)$. Coherent cohomology $H^*(X, \mathcal{O}_X)$ does not exactly compute the corner piece of the Hodge filtration, as would be an exact analogy with the formulation of Theorem 1.1, but it gives an upper bound. Indeed, by [14], Proposition 1.2, and [15], Proof of Theorem 1.1, one has a functorial surjective map $H^*(X, \mathcal{O}_X) \rightarrow \text{gr}_F^0 H^*(X, \mathbb{C})$.

The construction of these Witt vector cohomology spaces is given in section 2. If U is a separated k -scheme of finite type, $U \hookrightarrow X$ an open immersion in a proper k -scheme, and $\mathcal{I} \subset \mathcal{O}_X$ any coherent ideal such that $V(\mathcal{I}) = X \setminus U$, we define $W\mathcal{I} = \text{Ker}(W\mathcal{O}_X \rightarrow W(\mathcal{O}_X/\mathcal{I}))$, and we show that the cohomology spaces $H^*(X, W\mathcal{I}_K)$ actually depend only on U (Theorem 2.4). This results from an extension to Witt vectors of Deligne's results on the independence on the compactification for the construction of the $f_!$ functor for coherent sheaves [18].

When U varies, these spaces are contravariant functors with respect to proper maps, and covariant functors with respect to open immersions. In particular, they give rise to the usual long exact sequence relating the cohomologies of U , of an open subset $V \subset U$, and of the complement T of V in U . Thus, they can be viewed as providing a notion of Witt vector cohomology spaces with compact support for U . We define

$$H_c^*(U, W\mathcal{O}_{U,K}) := H^*(X, W\mathcal{I}_K).$$

Among other properties, we prove in section 3 that these cohomology spaces satisfy a particular case of cohomological descent (Theorem 3.2) which will be used in the proof of Theorem 1.1.

The construction of the canonical homomorphism between rigid and Witt vector cohomologies is given in section 4 (Theorem 4.5). First, we recall how to compute rigid cohomology for a proper k -scheme X when there exists a closed immersion of X in a smooth formal W -scheme \mathbb{P} , using the de Rham complex of \mathbb{P} with coefficients in an appropriate sheaf of $\mathcal{O}_{\mathbb{P}}$ -algebras $\mathcal{A}_{X,\mathbb{P}}$. When \mathbb{P} can be endowed with a lifting of the absolute Frobenius endomorphism of its special fibre, a simple construction (based on an idea of Illusie [20]) provides a functorial morphism from this de Rham complex to $W\mathcal{O}_{X,K}$. Using Čech resolutions, this morphism can still be defined in the general case as a morphism in the derived category of sheaves of K -vector spaces. Then we obtain (1.3) by taking the morphism induced on cohomology and restricting to the slope < 1 subspace. By means of simplicial resolutions based on de Jong's theorem, we prove in section 5 that it is an isomorphism. The proof proceeds by reduction to the case of proper and smooth schemes, using Theorem 3.2 and the descent properties of rigid cohomology proved by Chiarellotto and Tsuzuki ([10], [27], [28]).

We now list some applications of Theorem 1.1, which are developed in section 6. We first remark that it implies a vanishing theorem. If X is smooth projective, and $Y \subset X$ is a divisor so that $U = X \setminus Y$ is affine, then Serre vanishing says that $H^i(X, \mathcal{O}_X(-nY)) = 0$ for n large

and $i < \dim(X)$. If k has characteristic 0, then we can take $n = 1$ by Kodaira vanishing. One then has $H^i(X, \mathcal{O}_X(-Y)) = 0$ for $i < \dim(X)$. It is tempting to view the following corollary as an analogue when k has characteristic p :

Corollary 1.2. *Let X be a proper scheme defined over a perfect field k of characteristic $p > 0$. Let $\mathcal{I} \subset \mathcal{O}_X$ be a coherent ideal defining a closed subscheme $Y \subset X$ such that $U = X \setminus Y$ is affine, smooth and equidimensional of dimension n . Then*

$$H_c^i(U, W\mathcal{O}_{U,K}) = H^i(X, W\mathcal{I})_K = H^i(X, W\mathcal{I}^r)_K = 0$$

for all $i \neq n$ and all $r \geq 1$.

Indeed, when U is smooth and affine, $H_{\text{rig}}^i(U/K)$ can be identified with Monsky-Washnitzer cohomology [4], and therefore vanishes for $i > n = \dim(U)$. If moreover U is equidimensional, it follows by Poincaré duality [5] that $H_{\text{rig,c}}^i(U/K) = 0$ for $i < n$. So (1.3) implies that $H^i(X, W\mathcal{I})_K = 0$ for $i < n$. On the other hand, the closure \bar{U} of U in X has dimension n , and $H^i(X, W\mathcal{I})_K = H_c^i(U, W\mathcal{O}_{U,K})$ does not change up to canonical isomorphism if we replace X by \bar{U} . Therefore $H^i(X, W\mathcal{I})_K = 0$ for $i > n$.

When k is a finite field \mathbb{F}_q , with $q = p^a$, Theorem 1.1 implies a statement about zeta functions. By ([16], Théorème II), the Lefschetz trace formula provides an expression of the ζ -function of X as the alternating product

$$\zeta(X, t) = \prod_i P_i(X, t)^{(-1)^{i+1}},$$

where $P_i(X, t) = \det(1 - t\phi | H_{\text{rig,c}}^i(X/K))$, and $\phi = F^a$ denotes the \mathbb{F}_q -linear Frobenius endomorphism of X . On the other hand, we define

$$(1.4) \quad \begin{aligned} P_i^W(X, t) &= \det(1 - t\phi | H_c^i(X, W\mathcal{O}_{X,K})), \\ \zeta^W(X, t) &= \prod_i P_i^W(X, t)^{(-1)^{i+1}}. \end{aligned}$$

If we denote by $\zeta^{<1}(X, t)$ the “slope < 1 factor” of $\zeta(X, t)$ (cf. 6.1), we get formally from (1.3):

Corollary 1.3. *Let X be a separated scheme of finite type over a finite field. Then one has*

$$(1.5) \quad \zeta^{<1}(X, t) = \zeta^W(X, t).$$

This result can be used to prove congruences mod q on the number of \mathbb{F}_q -rational points of certain varieties. The following theorem answers a question of Serre, and was the initial motivation for this work. Recall that an effective divisor D on an abelian variety A is called a theta divisor if $\mathcal{O}_A(D)$ is ample and defines a principal polarization.

Theorem 1.4. *Let Θ, Θ' be two theta divisors on an abelian variety defined over a finite field \mathbb{F}_q . Then:*

$$(1.6) \quad |\Theta(\mathbb{F}_q)| \equiv |\Theta'(\mathbb{F}_q)| \pmod{q}.$$

Actually, Serre's original formulation predicts that, on an abelian variety defined over a field, the difference of the motives of Θ and Θ' is divisible by the Lefschetz motive. Our Theorem 1.4 answers the point counting consequence of it.

We also have more elementary point counting consequences.

Corollary 1.5 (Ax [1], Katz [21]). *Let $D_1, \dots, D_r \subset \mathbb{P}^n$ be hypersurfaces of degrees d_1, \dots, d_r , defined over the finite field \mathbb{F}_q . Assume that $\sum_j d_j \leq n$. Then*

$$(1.7) \quad |(D_1 \cap \dots \cap D_r)(\mathbb{F}_q)| \equiv 1 \pmod{q}.$$

We observe here that this congruence is the best approximation of the results of Ax and Katz that can be obtained using Witt vector cohomology, since this method only provides information on the slope < 1 factor of the zeta function. It would be worthwhile to have for higher slopes results similar to Theorem 1.1 which might give the full Ax-Katz congruences.

We also remark that Ax's theorem has a motivic proof [7], which of course is more powerful than this slope proof. Yet it is of interest to remark that Theorem 1.1 applies here as well. As for Theorem 1.4, it seems more difficult to formulate a motivic proof, as it would have to deal with non-effective motives (see the discussion in 6.6).

Let us mention finally the following general consequence of Theorem 1.1, which for example can be applied in the context of the work of Fu and Wan on mirror congruences for Calabi-Yau varieties ([29], [30]):

Corollary 1.6. *Let $f : X \rightarrow Y$ be a morphism between two proper \mathbb{F}_q -schemes. If f induces isomorphisms $f^* : H^i(Y, \mathcal{O}_Y) \xrightarrow{\sim} H^i(X, \mathcal{O}_X)$ for all $i \geq 0$, then*

$$(1.8) \quad |X(\mathbb{F}_q)| \equiv |Y(\mathbb{F}_q)| \pmod{q}.$$

Acknowledgements: It is a pleasure to thank Jean-Pierre Serre for his strong encouragement and his help. Theorem 1.4 was the main motivation for this work. We thank Luc Illusie for useful discussions. The

third named author thanks Eckart Viehweg for his interest and encouragement. Corollary 1.2 is inspired by the analogy on the relation between de Rham and coherent cohomology, which has been developed jointly with him.

Notations and conventions: Throughout this article, k denotes a perfect field of characteristic p , $W = W(k)$, $\sigma : W \rightarrow W$ the Frobenius automorphism of W , $K = \text{Frac}(W)$. The subscript K will denote tensorisation with K over W . We recall that, on any noetherian topological space, taking cohomology commutes with tensorisation by \mathbb{Q} . Therefore the subscript K will be moved inside or outside cohomology or direct images without further justification.

We denote by $D^b(K)$ (resp. $D^b(X, K)$) the derived category of bounded complexes of K -vector spaces (resp. complexes of sheaves of K -vector spaces on a topological space X).

All formal schemes considered in this article are W -formal schemes for the p -adic topology.

2. WITT VECTOR COHOMOLOGY WITH COMPACT SUPPORTS

We give in this section some properties of Witt vector cohomology which are a strong indication of its topological nature, and will be used later in our proof of Theorem 1.1. In particular, we show how to attach Witt vector cohomology spaces with compact supports to any separated k -scheme of finite type.

If X is a scheme, \mathcal{A} a sheaf of rings on X , and $n \geq 1$, we denote by $W_n\mathcal{A}$, or $W_n(\mathcal{A})$ if confusion may arise, the sheaf of Witt vectors of length n with coefficients in \mathcal{A} , and by $W\mathcal{A} = \varprojlim_n W_n\mathcal{A}$, or $W(\mathcal{A})$, the sheaf of Witt vectors of infinite length. If $\mathcal{I} \subset \mathcal{A}$ is an ideal, we denote by $W_n\mathcal{I} = \text{Ker}(W_n\mathcal{A} \rightarrow W_n(\mathcal{A}/\mathcal{I}))$, or $W_n(\mathcal{I})$, the sheaf of Witt vectors $(a_0, a_1, \dots, a_{n-1})$ such that a_i is a section of \mathcal{I} for all i , and we define similarly $W\mathcal{I}$. Note that, when \mathcal{I} is quasi-coherent, the canonical morphism $W\mathcal{I} \rightarrow \mathbb{R}\varprojlim_n W_n\mathcal{I}$ is an isomorphism, as $H^1(U, W_n\mathcal{I}) = 0$ for any affine open subset $U \subset X$ and any n , and the projective system $\Gamma(U, W_n\mathcal{I})$ has surjective transition maps.

For any X , any sheaf of rings \mathcal{A} on X , and any ideal $\mathcal{I} \subset \mathcal{A}$, we use the notations $\mathbb{R}\Gamma(X, W\mathcal{I})$ and $H^*(X, W\mathcal{I})$ to denote the Zariski cohomology of the sheaf $W\mathcal{I}$. Thus, when \mathcal{I} is quasi-coherent, the canonical morphism

$$\mathbb{R}\Gamma(X, W\mathcal{I}) \rightarrow \mathbb{R}\varprojlim_n \mathbb{R}\Gamma(X, W_n\mathcal{I})$$

is an isomorphism. When X is a proper k -scheme and $\mathcal{I} \subset \mathcal{O}_X$ is a coherent ideal, the cohomology modules $H^*(X, W_n\mathcal{I})$ are artinian W -modules; then it follows from the Mittag-Leffler criterium that the morphism

$$H^*(X, W\mathcal{I}) \rightarrow \varprojlim_n H^*(X, W_n\mathcal{I})$$

is an isomorphism.

We will shorten notations by writing $W\mathcal{O}_{X,K}$, $W\mathcal{I}_K$ for $(W\mathcal{O}_X)_K$, $(W\mathcal{I})_K$. We recall again that, when X is noetherian, there is a canonical isomorphism

$$\mathbb{R}\Gamma(X, W\mathcal{I})_K \xrightarrow{\sim} \mathbb{R}\Gamma(X, W\mathcal{I}_K).$$

In this article, we will be particularly interested in the K -vector spaces $H^*(X, W\mathcal{O}_{X,K})$, and in their generalizations $H^*(X, W\mathcal{I}_K)$. Our main observation is that, in contrast to the W -modules $H^*(X, W\mathcal{O}_X)$, which are sensitive to nilpotent sections of \mathcal{O}_X , they behave like a topological cohomology theory for separated k -schemes of finite type. The following easy proposition is somehow a key point.

Proposition 2.1. *Let X be a k -scheme of finite type.*

(i) *The canonical homomorphism*

$$W\mathcal{O}_{X,K} \rightarrow W\mathcal{O}_{X_{\text{red}},K}$$

is an isomorphism, and induces a functorial isomorphism

$$H^*(X, W\mathcal{O}_{X,K}) \xrightarrow{\sim} H^*(X_{\text{red}}, W\mathcal{O}_{X_{\text{red}},K}),$$

compatible with the action of F and V .

(ii) *Let $\mathcal{I}, \mathcal{J} \subset \mathcal{O}_X$ be coherent ideals, and assume $\sqrt{\mathcal{I}} = \sqrt{\mathcal{J}}$, i.e. there exists $N \geq 1$ such that $\mathcal{I}^N \subset \mathcal{J}$ and $\mathcal{J}^N \subset \mathcal{I}$. Then there is a canonical identification*

$$W\mathcal{I}_K \cong W\mathcal{J}_K,$$

inducing a functorial isomorphism

$$H^*(X, W\mathcal{I}_K) \cong H^*(X, W\mathcal{J}_K),$$

compatible with the action of F and V .

Proof. Let $\mathcal{N} = \text{Ker}(\mathcal{O}_X \rightarrow \mathcal{O}_{X_{\text{red}}})$. To prove the first claim of assertion (i), it suffices to show that $W\mathcal{N}_K = 0$. But the action of p is invertible, and $p = VF = FV$, so it suffices to show F is nilpotent. This is clear since F acts on $W\mathcal{N}$ by raising coordinates to the p -th power. Taking cohomology, the second claim follows.

Assertion (ii) follows from assertion (i) applied to $\mathcal{O}_X/\mathcal{I}$ and $\mathcal{O}_X/\mathcal{J}$. \square

We now prove the existence of Mayer-Vietoris exact sequences for Witt vector cohomology.

Proposition 2.2. *Let X be a k -scheme of finite type, $X_1, X_2 \subset X$ two closed subschemes such that $X = X_1 \cup X_2$, and $Z = X_1 \cap X_2$. There is an exact sequence*

$$(2.1) \quad 0 \rightarrow W\mathcal{O}_{X,K} \rightarrow W\mathcal{O}_{X_1,K} \oplus W\mathcal{O}_{X_2,K} \rightarrow W\mathcal{O}_{Z,K} \rightarrow 0,$$

providing a Mayer-Vietoris long exact sequence

$$\begin{aligned} \cdots \rightarrow H^i(X, W\mathcal{O}_{X,K}) &\rightarrow H^i(X_1, W\mathcal{O}_{X_1,K}) \oplus H^i(X_2, W\mathcal{O}_{X_2,K}) \\ &\rightarrow H^i(Z, W\mathcal{O}_{Z,K}) \rightarrow H^{i+1}(X, W\mathcal{O}_{X,K}) \rightarrow \cdots \end{aligned}$$

Proof. Let $\mathcal{I}_1, \mathcal{I}_2$ be the ideals of \mathcal{O}_X defining X_1 and X_2 . Thanks to 2.1, we may assume that $\mathcal{O}_Z = \mathcal{O}_X/(\mathcal{I}_1 + \mathcal{I}_2)$. It is easy to check that $W(\mathcal{I}_1 + \mathcal{I}_2) = W(\mathcal{I}_1) + W(\mathcal{I}_2)$. From the exact sequence

$$0 \rightarrow \mathcal{O}_X/(\mathcal{I}_1 \cap \mathcal{I}_2) \rightarrow \mathcal{O}_X/\mathcal{I}_1 \oplus \mathcal{O}_X/\mathcal{I}_2 \rightarrow \mathcal{O}_X/(\mathcal{I}_1 + \mathcal{I}_2) \rightarrow 0,$$

we can then deduce an exact sequence

$$0 \rightarrow W(\mathcal{O}_X/(\mathcal{I}_1 \cap \mathcal{I}_2)) \rightarrow W\mathcal{O}_{X_1} \oplus W\mathcal{O}_{X_2} \rightarrow W\mathcal{O}_Z \rightarrow 0.$$

Since $X = X_1 \cup X_2$, $\mathcal{I}_1 \cap \mathcal{I}_2$ is a nilpotent ideal, and Proposition 2.1 implies that $W\mathcal{O}_{X,K} \xrightarrow{\sim} W(\mathcal{O}_X/(\mathcal{I}_1 \cap \mathcal{I}_2))_K$. This gives the short exact sequence of the statement. The long one follows by taking cohomology. \square

Corollary 2.3. *Let X be a k -scheme of finite type, $X_1, \dots, X_r \subset X$ closed subschemes such that $X = X_1 \cup \cdots \cup X_r$. For each sequence $1 \leq i_0 < \cdots < i_n \leq r$, let $X_{i_0, \dots, i_n} = X_{i_0} \cap \cdots \cap X_{i_n}$. Then the sequence*

$$(2.2) \quad 0 \rightarrow W\mathcal{O}_{X,K} \rightarrow \prod_{i=1}^r W\mathcal{O}_{X_i,K} \rightarrow \cdots \rightarrow W\mathcal{O}_{X_1, \dots, r, K} \rightarrow 0$$

is exact.

Proof. The statement being true for $r = 2$, we proceed by induction on r . Let $X' = X_2 \cup \cdots \cup X_r$. Up to a shift, the complex (2.2) is the cone of the morphism of complexes

$$(2.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & W\mathcal{O}_{X,K} & \longrightarrow & \prod_{i=2}^r W\mathcal{O}_{X_i,K} & \longrightarrow & \cdots \longrightarrow W\mathcal{O}_{X_2, \dots, r, K} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & W\mathcal{O}_{X_1,K} & \longrightarrow & \prod_{i=2}^r W\mathcal{O}_{X_{1,i},K} & \longrightarrow & \cdots \longrightarrow W\mathcal{O}_{X_1, \dots, r, K} \longrightarrow 0. \end{array}$$

Thus it suffices to prove that this morphism is a quasi-isomorphism. The induction hypothesis implies that the canonical morphism from

the complex $0 \rightarrow W\mathcal{O}_{X,K} \rightarrow W\mathcal{O}_{X',K} \rightarrow 0$ (resp. $0 \rightarrow W\mathcal{O}_{X_1,K} \rightarrow W\mathcal{O}_{X_1 \cap X',K} \rightarrow 0$) to the upper line (resp. lower line) of (2.3) is a quasi-isomorphism. Therefore it suffices to prove that the morphism of complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & W\mathcal{O}_{X,K} & \longrightarrow & W\mathcal{O}_{X',K} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & W\mathcal{O}_{X_1,K} & \longrightarrow & W\mathcal{O}_{X_1 \cap X',K} & \longrightarrow & 0 \end{array}$$

is a quasi-isomorphism, and this follows from Proposition 2.2. \square

Our next theorem is the main result which allows to use the groups $H^*(X, W\mathcal{I}_K)$ to define a Witt vector cohomology with compact supports for separated k -schemes of finite type. Its starting point is Deligne's "independence on the compactification" result in the construction of the $f_!$ functor for coherent sheaves [18, App., Prop. 5].

Theorem 2.4. *Let $f : X' \rightarrow X$ be a proper morphism between two k -schemes of finite type, and let $\mathcal{I} \subset \mathcal{O}_X$ be a coherent ideal, $\mathcal{I}' = \mathcal{I}\mathcal{O}_{X'}$, $Y = V(\mathcal{I})$, $U = X \setminus Y$, $U' = f^{-1}(U)$.*

(i) *If f induces a finite morphism $U' \rightarrow U$, then, for all $q \geq 1$,*

$$(2.4) \quad R^q f_*(W\mathcal{I}'_K) = 0.$$

(ii) *If f induces an isomorphism $U' \xrightarrow{\sim} U$, the canonical morphism*

$$(2.5) \quad W\mathcal{I}_K \rightarrow f_*(W\mathcal{I}'_K)$$

is an isomorphism, and induces an isomorphism

$$(2.6) \quad \mathbb{R}\Gamma(X, W\mathcal{I}_K) \xrightarrow{\sim} \mathbb{R}\Gamma(X', W\mathcal{I}'_K).$$

The proof will use the following general lemma:

Lemma 2.5. *Let $I \subset A$ be an ideal in a ring A . For all integers $n \geq 2, r, s \in \mathbb{N}$, define*

$$W_n^{r,s}(I) = \{(a_0, a_1, \dots) \in W_n(A) \mid a_0 \in I^r, a_i \in I^s \text{ for all } i \geq 1\}.$$

(i) *If $s \leq pr$, the subset $W_n^{r,s}(I)$ is an ideal of $W_n(A)$, which sits in the short exact sequence of abelian groups*

$$(2.7) \quad 0 \rightarrow W_{n-1}(I^s) \xrightarrow{V} W_n^{r,s}(I) \xrightarrow{R^{n-1}} I^r \rightarrow 0.$$

(ii) If $r \leq s \leq pr$, the sequence (2.7) sits in a commutative diagram of short exact sequences

$$(2.8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & W_{n-1}(I^s) & \xrightarrow{V} & W_n(I^s) & \xrightarrow{R^{n-1}} & I^s \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & W_{n-1}(I^s) & \xrightarrow{V} & W_n^{r,s}(I) & \xrightarrow{R^{n-1}} & I^r \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & W_{n-1}(I^r) & \xrightarrow{V} & W_n(I^r) & \xrightarrow{R^{n-1}} & I^r \longrightarrow 0 \end{array}$$

where the vertical arrows are the natural inclusions.

Proof. Assume first that $s \leq pr$. The subset $W_n^{r,s}(I)$ can be described as the set of Witt vectors of the form $\underline{a} + V(b)$, where $a \in I^r$, $b \in W_{n-1}(I^s)$, and \underline{a} denotes the Teichmüller representative of a . To prove that it is an additive subgroup, it suffices to verify that if $a, a' \in I^r$, then $\underline{a} + \underline{a}' = \underline{a + a'} + V(c)$, with $c \in W_{n-1}(I^s)$. If $S_i(X_0, \dots, X_i, Y_0, \dots, Y_i)$ are the universal polynomials defining the addition in $W_n(A)$, it is easy to check that $S_i(X_0, 0, \dots, Y_0, 0, \dots)$ is a homogeneous polynomial of degree p^i in $\mathbb{Z}[X_0, Y_0]$ [26, II §6]. Since $pr \geq s$, the claim follows. Using again the condition $pr \geq s$, the fact that $W_n^{r,s}(I)$ satisfies the multiplicativity property of an ideal follows from the relations

$$\underline{a}(x_0, x_1, x_2, \dots) = (ax_0, a^p x_1, a^{p^2} x_2, \dots), \quad V(b)x = V(bF(x)).$$

If we assume in addition that $r \leq s$, then the vertical inclusions of the diagram are defined, and its commutativity is obvious. \square

We will use repeatedly the following elementary remark.

Lemma 2.6. *Let \mathcal{C} be an abelian category, and let*

$$(2.9) \quad \begin{array}{ccccc} & & E & & \\ & & \downarrow v & \searrow u'' & \\ G' & \xrightarrow{V} & G & \xrightarrow{R} & G'' \\ & \searrow u' & \downarrow w & & \\ & & F & & \end{array}$$

be a commutative diagram of morphisms of \mathcal{C} such that the horizontal sequence is exact. If $u' = 0$ and $u'' = 0$, then $w \circ v = 0$.

Proof. Exercise. \square

Lemma 2.7. (i) *Under the assumptions of Theorem 2.4, (i) there exists an integer $a \geq 0$ such that, for all $q \geq 1$, all $n \geq 1$ and all $r \geq 0$, the canonical morphism*

$$(2.10) \quad R^q f_*(W_n(\mathcal{I}^{r+a})) \rightarrow R^q f_*(W_n(\mathcal{I}^r))$$

is the zero morphism.

(ii) *Assume that f induces an isomorphism $U' \xrightarrow{\sim} U$, and define*

$$\mathcal{K}_n^r = \text{Ker}(W_n(\mathcal{I}^r) \rightarrow f_*(W_n(\mathcal{I}^r))),$$

$$\mathcal{C}_n^r = \text{Coker}(W_n(\mathcal{I}^r) \rightarrow f_*(W_n(\mathcal{I}^r))).$$

Then there exists an integer $a \geq 0$ such that, for all $n \geq 1$ and all $r \geq 0$, the canonical morphisms

$$(2.11) \quad \mathcal{K}_n^{r+a} \rightarrow \mathcal{K}_n^r, \quad \mathcal{C}_n^{r+a} \rightarrow \mathcal{C}_n^r,$$

are the zero morphisms.

Proof. We first prove assertion (i). We may fix $q \geq 1$, since $R^q f_* = 0$ for q big enough. When $n = 1$, we can apply Deligne's result [18, App., Prop. 5] to $\mathcal{O}_{X'}$, and we obtain an integer $b \geq 0$ such that, for all $r \geq 0$, the canonical morphism

$$u_1 : R^q f_*(\mathcal{I}^{r+b}) \rightarrow R^q f_*(\mathcal{I}^r)$$

is 0 (note that, for this result, Deligne's argument only uses that f is finite above U). Let us prove by induction on n that, for all $n \geq 1$ and all $r \geq b$, the canonical morphism

$$u_n : R^q f_*(W_n(\mathcal{I}^{r+b})) \rightarrow R^q f_*(W_n(\mathcal{I}^r))$$

is also 0.

The condition $r \geq b$ implies that the couple $(r, r+b)$ is such that $r \leq r+b \leq pr$. Therefore, we may use Lemma 2.5 to define ideals $W_n^{r, r+b}(\mathcal{I}') \subset W_n(\mathcal{O}_{X'})$, and, for $n \geq 2$, we obtain a commutative diagram (2.8) relative to \mathcal{I}' and $(r, r+b)$. Since the middle row of (2.8) is a short exact sequence, the diagram

$$\begin{array}{ccccc} & & R^q f_*(W_n(\mathcal{I}^{r+b})) & \xrightarrow{R^{n-1}} & R^q f_*(\mathcal{I}^{r+b}) \\ & & \downarrow & & \downarrow u_1 \\ R^q f_*(W_{n-1}(\mathcal{I}^{r+b})) & \xrightarrow{V} & R^q f_*(W_n^{r, r+b}(\mathcal{I}')) & \xrightarrow{R^{n-1}} & R^q f_*(\mathcal{I}^r) \\ & & \downarrow & & \downarrow \\ & & R^q f_*(W_{n-1}(\mathcal{I}^r)) & \xrightarrow{V} & R^q f_*(W_n(\mathcal{I}^r)) \end{array}$$

obtained by applying the functor $R^q f_*$ to (2.8) has an exact middle row. As $u_1 = 0$, and the composition of the middle vertical arrows is

u_n , Lemma 2.6 shows by induction that $u_n = 0$ for all $n \geq 1$. If we set $a = 2b$, then assertion (i) holds.

Assertion (ii) can be proved by the same method. Let $\mathcal{R} = \bigoplus_{r \geq 0} \mathcal{I}^r$. It follows from [17, 3.3.1] that the quasi-coherent graded \mathcal{R} -modules $\mathcal{K}_1 = \bigoplus_{r \geq 0} \mathcal{K}_1^r$ and $\mathcal{C}_1 = \bigoplus_{r \geq 0} \mathcal{C}_1^r$ are finitely generated. As the restriction of f to U' is an isomorphism, they are supported in Y . Therefore, there exists an integer m such that $\mathcal{I}^m \mathcal{K}_1^r = \mathcal{I}^m \mathcal{C}_1^r = 0$ for all $r \geq 0$. Moreover, there exists an integer d such that, for all $r \geq d$ the canonical morphisms

$$\mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{K}_1^r \rightarrow \mathcal{K}_1^{r+1}, \quad \mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{C}_1^r \rightarrow \mathcal{C}_1^{r+1},$$

are surjective. It follows that the morphisms

$$\mathcal{K}_1^{r+m} \rightarrow \mathcal{K}_1^r, \quad \mathcal{C}_1^{r+m} \rightarrow \mathcal{C}_1^r$$

are 0 for $r \geq d$. Replacing m by $b = d + m$, the corresponding morphisms are 0 for all $r \geq 0$, which proves assertion (ii) when $n = 1$.

As $\mathcal{K}_1^r \subset \mathcal{O}_X$ for all r , this implies that $\mathcal{K}_1^r = 0$ for $r \geq b$. Thanks to the exact sequences

$$0 \rightarrow \mathcal{K}_{n-1}^r \xrightarrow{V} \mathcal{K}_n^r \xrightarrow{R^{n-1}} \mathcal{K}_1^r,$$

it follows that $\mathcal{K}_n^r = 0$ for all $n \geq 1$ and all $r \geq b$. Thus assertion (ii) holds for the modules \mathcal{K}_n^r , with $a = b$.

To prove it for the modules \mathcal{C}_n^r , we introduce for $r \geq b$ the modules

$$\mathcal{C}_n^{r,r+b} = \text{Coker}(W_n^{r,r+b}(\mathcal{I}) \rightarrow f_*(W_n^{r,r+b}(\mathcal{I}'))).$$

The snake lemma applied to the diagrams

$$\begin{array}{ccccccc} 0 & \longrightarrow & W_{n-1}(\mathcal{I}^{r+b}) & \xrightarrow{V} & W_n^{r,r+b}(\mathcal{I}) & \xrightarrow{R^{n-1}} & \mathcal{I}^r \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & f_*(W_{n-1}(\mathcal{I}^{r+b})) & \xrightarrow{V} & f_*(W_n^{r,r+b}(\mathcal{I}')) & \xrightarrow{R^{n-1}} & f_*(\mathcal{I}^r) \end{array}$$

gives exact sequences

$$\mathcal{C}_{n-1}^{r+b} \xrightarrow{V} \mathcal{C}_n^{r,r+b} \xrightarrow{R^{n-1}} \mathcal{C}_1^r.$$

The functoriality of the diagrams (2.8) imply that these exact sequences sit in commutative diagrams

$$\begin{array}{ccccc}
 & & \mathcal{C}_n^{r+b} & \xrightarrow{R^{n-1}} & \mathcal{C}_1^{r+b} \\
 & & \downarrow & & \downarrow u_1 \\
 \mathcal{C}_{n-1}^{r+b} & \xrightarrow{V} & \mathcal{C}_n^{r,r+b} & \xrightarrow{R^{n-1}} & \mathcal{C}_1^r \\
 \downarrow u_{n-1} & & \downarrow & & \\
 \mathcal{C}_{n-1}^r & \xrightarrow{V} & \mathcal{C}_n^r & & ,
 \end{array}$$

where the morphisms u_i are the canonical morphisms, and the composition of the middle vertical arrows is u_n . As $u_1 = 0$, Lemma 2.6 implies by induction that $u_n = 0$ for all n . This proves assertion (ii) for \mathcal{C}_n^r , with $a = 2b$. \square

2.8. Proof of Theorem 2.4. Under the assumptions of 2.4, let $q \geq 1$ be an integer, and let a be an integer satisfying the conclusion of Lemma 2.7 (i) for the family of sheaves $R^q f_*(W_n(\mathcal{I}^r))$, for all $n \geq 1$ and $r \geq 0$. Let c be such that $p^c > a$. Since the Frobenius map $F^c : R^q f_*(W_n(\mathcal{I}^r)) \rightarrow R^q f_*(W_n(\mathcal{I}^r))$ factors through $R^q f_*(W_n(\mathcal{I}^{p^c r}))$, it follows from Lemma 2.7 that F^c acts by zero on $R^q f_*(W_n(\mathcal{I}^r))$, for all $n \geq 1$ and all $r \geq 1$. Therefore, for all $r \geq 1$, F^c acts by zero on $\mathbb{R} \varprojlim_n (R^q f_*(W_n(\mathcal{I}^r)))$.

In particular, F^c acts by zero on the sheaves $\varprojlim_n R^q f_*(W_n \mathcal{I}')$ and $R^1 \varprojlim_n R^q f_*(W_n \mathcal{I}')$. As the inverse system $(W_n \mathcal{I}')_{n \geq 1}$ has surjective transition maps, and terms with vanishing cohomology on affine open subsets, it is \varprojlim_n -acyclic, and we obtain

$$\mathbb{R} f_*(W \mathcal{I}') = \mathbb{R} f_*(\mathbb{R} \varprojlim_n W_n \mathcal{I}') \cong \mathbb{R} \varprojlim_n \mathbb{R} f_*(W_n \mathcal{I}').$$

This isomorphism provides a biregular spectral sequence

$$E_2^{i,j} = R^i \varprojlim_n R^j f_*(W_n \mathcal{I}') \Rightarrow R^{i+j} f_*(W \mathcal{I}'),$$

in which the filtration of the terms $R^{i+j} f_*(W \mathcal{I}')$ is of length ≤ 2 since the functors $R^i \varprojlim_n$ are zero for $i \geq 2$. As F^c acts by zero on the terms $E_2^{i,j}$ for $j \geq 1$, F^{2c} acts by zero on the terms $R^{i+j} f_*(W \mathcal{I}')$ for $i+j \geq 2$.

On the other hand, the term $E_2^{1,0} = R^1 \varprojlim_n f_*(W_n \mathcal{I}')$ is 0, because the morphisms $f_*(W_{n+1} \mathcal{I}') \rightarrow f_*(W_n \mathcal{I}')$ are surjective (since $f_*(W_n \mathcal{I}') \cong W_n(f_*(\mathcal{I}'))$), and the cohomology of the terms $f_*(W_n \mathcal{I}')$ vanishes on any open affine subset. Therefore, F^c acts by zero on $R^1 f_*(W \mathcal{I}')$.

Thus the action of F on $R^q f_*(W\mathcal{I}')$ is nilpotent for all $q \geq 1$. Since this action becomes an isomorphism after tensorisation with K , we obtain that $R^q f_*(W\mathcal{I}'_K) = R^q f_*(W\mathcal{I}')_K = 0$ for all $q \geq 1$, which proves assertion (i) of Theorem 2.4.

Let us assume now that $f : U' \rightarrow U$ is an isomorphism. It follows from Lemma 2.7 (ii) that there exists an integer a such that, for all $n \geq 1$ and all $r \geq 0$, the morphisms $\mathcal{K}_n^{r+a} \rightarrow \mathcal{K}_n^r$ and $\mathcal{C}_n^{r+a} \rightarrow \mathcal{C}_n^r$ are zero. Taking c such that $p^c > a$, it follows that F^c acts by zero on \mathcal{K}_n^r and \mathcal{C}_n^r for all n and all r . Therefore, F^c acts by zero on $\varprojlim_n \mathcal{K}_n^1$ and $\varprojlim_n \mathcal{C}_n^1$. It is easy to check that these two inverse systems and the inverse system $\text{Im}(W_n\mathcal{I} \rightarrow f_*(W_n\mathcal{I}'))$ all have surjective transition maps. As their terms have vanishing cohomology on affine open subsets, they are \varprojlim -acyclic, and we obtain

$$\begin{aligned} \text{Ker}(W\mathcal{I} \rightarrow f_*(W\mathcal{I}')) &= \varprojlim_n \mathcal{K}_n^1, \\ \text{Coker}(W\mathcal{I} \rightarrow f_*(W\mathcal{I}')) &= \varprojlim_n \mathcal{C}_n^1. \end{aligned}$$

After tensorisation with K , F becomes an isomorphism on $\varprojlim_n \mathcal{K}_n^1$ and $\varprojlim_n \mathcal{C}_n^1$, and assertion (ii) of Theorem 2.4 follows. \square

2.9. We now observe that the previous results imply that the cohomology spaces $H^*(X, W\mathcal{I}_K)$ only depend on the k -scheme U , and have the same functoriality properties with respect to U than cohomology with compact supports.

Indeed, suppose U is fixed, and let $U \hookrightarrow X_1, U \hookrightarrow X_2$ be two open immersions into proper k -schemes. Let $X \subset X_1 \times_k X_2$ be the scheme theoretic closure of U embedded diagonally into $X_1 \times_k X_2$. The two projections induce proper maps $p_1 : X \rightarrow X_1, p_2 : X \rightarrow X_2$. If $\mathcal{I} \subset \mathcal{O}_X, \mathcal{I}_1 \subset \mathcal{O}_{X_1}, \mathcal{I}_2 \subset \mathcal{O}_{X_2}$ are the ideals defining $Y = (X \setminus U)_{\text{red}}, Y_1 = (X_1 \setminus U)_{\text{red}}$ and $Y_2 = (X_2 \setminus U)_{\text{red}}$, we deduce from 2.1 and 2.4 that the homomorphisms

$$\mathbb{R}\Gamma(X_2, W\mathcal{I}_{2,K}) \xrightarrow{p_2^*} \mathbb{R}\Gamma(X, W\mathcal{I}_K) \xleftarrow{p_1^*} \mathbb{R}\Gamma(X_1, W\mathcal{I}_{1,K})$$

are isomorphisms. If we define

$$\varepsilon_{12} = p_1^{*-1} \circ p_2^* : \mathbb{R}\Gamma(X_2, W\mathcal{I}_{2,K}) \xrightarrow{\sim} \mathbb{R}\Gamma(X_1, W\mathcal{I}_{1,K}),$$

it is easy to check that the isomorphisms ε_{ij} satisfy the transitivity condition for a third open immersion $U \hookrightarrow X_3$ into a proper k -scheme. Therefore, they provide canonical identifications between the cohomology complexes $\mathbb{R}\Gamma(X, W\mathcal{I}_K)$ defined by various open immersions of U into proper k -schemes X .

Assume U is a separated k -scheme of finite type. Since, by Nagata's theorem, there exists a proper k -scheme X and an open immersion $U \hookrightarrow X$, this independence property allows to define the *Witt vector cohomology with compact supports* of U by setting

$$(2.12) \quad \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K}) := \mathbb{R}\Gamma(X, W\mathcal{I}_K) \cong \mathbb{R}\Gamma(X, W\mathcal{I})_K,$$

$$(2.13) \quad H_c^*(U, W\mathcal{O}_{U,K}) := H^*(X, W\mathcal{I}_K) \cong H^*(X, W\mathcal{I})_K,$$

where $\mathcal{I} \subset \mathcal{O}_X$ is any coherent ideal defining the closed subset $X \setminus U$. As the restriction of $W\mathcal{I}$ to U is $W\mathcal{O}_U$, there is a canonical morphism $\mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K}) \rightarrow \mathbb{R}\Gamma(U, W\mathcal{O}_{U,K})$, which is an isomorphism when U is proper.

These cohomology groups have the following functoriality properties:

(i) They are contravariant with respect to proper maps.

Let $f : U' \rightarrow U$ be a proper k -morphism of separated k -schemes, and let $U' \hookrightarrow X', U \hookrightarrow X$ be open immersions into proper k -schemes. Replacing if necessary X' by the scheme theoretic closure of the graph of f in $X' \times_k X$, we may assume that there exists a k -morphism $g : X' \rightarrow X$ extending f , and that U' is dense in X' . As f is proper, it follows that $U' = g^{-1}(U)$. If $\mathcal{I} \subset \mathcal{O}_X$ is any coherent ideal such that $V(\mathcal{I}) = X \setminus U$, then $\mathcal{I}' = \mathcal{I}\mathcal{O}_{X'}$ is such that $V(\mathcal{I}') = X' \setminus U'$, and we can define the homomorphism

$$f^* : \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K}) \rightarrow \mathbb{R}\Gamma_c(U', W\mathcal{O}_{U',K})$$

as being

$$g^* : \mathbb{R}\Gamma(X, W\mathcal{I}_K) \rightarrow \mathbb{R}\Gamma(X', W\mathcal{I}'_K).$$

We leave as an exercise to check that, up to canonical isomorphism, f^* does not depend on the choices.

(ii) They are covariant with respect to open immersions.

Let $j : V \hookrightarrow U$ be an open immersion, let $U \hookrightarrow X$ be an open immersion into a proper k -scheme, and let $\mathcal{I}, \mathcal{J} \subset \mathcal{O}_X$ be the ideals of $Y = (X \setminus U)_{\text{red}}$ and $Z = (X \setminus V)_{\text{red}}$. Then the homomorphism

$$j_* : \mathbb{R}\Gamma_c(V, W\mathcal{O}_{V,K}) \rightarrow \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K})$$

is defined as being

$$\mathbb{R}\Gamma(X, W\mathcal{J}_K) \rightarrow \mathbb{R}\Gamma(X, W\mathcal{I}_K).$$

If $T = U \setminus V = Z \cap U$, then T is open in the proper k -scheme Z , and the ideal $\mathcal{I}/\mathcal{J} \subset \mathcal{O}_Z$ defines the complement of T in Z . Thus, the usual distinguished triangle

$$(2.14) \quad \mathbb{R}\Gamma_c(V, W\mathcal{O}_{V,K}) \rightarrow \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K}) \rightarrow \mathbb{R}\Gamma_c(T, W\mathcal{O}_{T,K}) \xrightarrow{+1}$$

is obtained by tensoring the short exact sequence

$$0 \rightarrow W\mathcal{J} \rightarrow W\mathcal{I} \rightarrow W(\mathcal{I}/\mathcal{J}) \rightarrow 0,$$

with K , and taking its cohomology on X .

Proposition 2.10. *For any separated k -scheme of finite type U , the cohomology spaces $H_c^*(U, W\mathcal{O}_{U,K})$ are finite dimensional K -vector spaces, on which the Frobenius endomorphism has slopes in $[0, 1[$.*

Proof. It suffices to prove the statement when U is a proper k -scheme X . Then the statement reduces to the finiteness of the usual cohomology spaces $H^*(X, W\mathcal{O}_X)_K$. This is a well-known result (cf. [6, III, Th. 2.2], which is valid for $W\mathcal{O}_X$ without the smoothness assumption). For the sake of completeness, we give a proof here.

Write $M := H^i(X, W\mathcal{O}_X)$. As X is proper, the groups $H^i(X, W_n\mathcal{O}_X)$ satisfy the Mittag-Leffler condition, so that the homomorphism

$$H^i(X, W\mathcal{O}_X) \rightarrow \varprojlim_n H^i(X, W_n\mathcal{O}_X)$$

is an isomorphism. Therefore, the Verschiebung endomorphism endows M with a structure of module over the non necessarily commutative ring $R := W_\sigma[[V]]$, where the index σ refers to the commutation rules $aV = V\sigma(a)$ for $a \in W$. Then M is separated and complete for the V -adic topology. On the other hand, $M/VM \hookrightarrow H^i(X, \mathcal{O}_X)$ is a finite dimensional k -vector space. Since M/VM is finitely generated over W , it follows that M is finitely generated over R . Moreover, the fact that M/VM has finite length implies that M is a torsion R -module. Thus, there exists a finite number of non zero elements $a_i(V) = a_{i,0} + a_{i,1}V + a_{i,2}V^2 + \dots \in R, i = 1, \dots, r$, and a surjection

$$(2.15) \quad \bigoplus_{i=1}^r R/Ra_i(V) \twoheadrightarrow M.$$

Fix some $i, 1 \leq i \leq r$. We are interested in the module structure after inverting p , so we may assume that $a_{i,\ell} \in W(k)^\times$ for some ℓ . Let ℓ be minimal, so $p|a_{i,j}, 0 \leq j < \ell$. Even if the ring R is not commutative, we can get a factorization

$$\sum_j a_{i,j}V^j = \left(c_{i,0} + c_{i,1}V + \dots\right) \left(b_{i,0} + \dots + b_{i,\ell-1}V^{\ell-1} + V^\ell\right); \quad p|b_{i,j}.$$

To see this, we factor

$$\sum_j a_{i,j}V^j \equiv \left(c_{i,0}^{(s)} + c_{i,1}^{(s)}V + \dots\right) \left(b_{i,0}^{(s)} + \dots + b_{i,\ell-1}^{(s)}V^{\ell-1} + V^\ell\right) \pmod{p^{s+1}},$$

where $b_{i,j}^{(s+1)}, c_{i,j}^{(s+1)} \equiv b_{i,j}^{(s)}, c_{i,j}^{(s)} \pmod{p^{s+1}}$, starting with $b_{i,j}^{(0)} = 0$ and $c_{i,j}^{(0)} = a_{i,j+\ell}$. Note that $c_{i,0}^{(s)} \in W(k)^\times$. Details are standard and are left for the reader. Writing $b_i(V) = b_{i,0} + b_{i,1}V + \dots + V^\ell$ with $b_{i,j} = \lim b_{i,j}^{(s)}$, it follows that $R/Ra_i(V) \cong R/Rb_i(V) \cong W(k)^{\oplus \ell}$. The finiteness now follows from (2.15).

As $H^*(X, W\mathcal{O}_X)$ is a finitely generated W -module modulo torsion, the slopes of the Frobenius endomorphism are positive. On the other hand, the existence of a Verschiebung endomorphism V such that $FV = VF = p$ implies that all slopes are ≤ 1 . As V is topologically nilpotent, there cannot be any non zero element of slope 1 in $H^*(X, W\mathcal{O}_{X,K})$, and all its slopes belong to $[0, 1[$. \square

3. A DESCENT THEOREM

We will need simplicial resolutions based on de Jong's fundamental result on alterations. In this section, we briefly recall some related definitions, and how to construct such resolutions. We then prove for Witt vector cohomology with compact supports a particular case of étale cohomological descent which will be one of the main ingredients in our proof of Theorem 1.1.

If X is a scheme, and $n \geq -1$ an integer, we denote as usual by sk_n^X the truncation functor from the category of simplicial X -schemes to the category of n -truncated simplicial X -schemes, and by cosk_n^X its right adjoint [12, 5.1].

Definition 3.1. Let X be a reduced k -scheme of finite type, and let X_\bullet be a simplicial k -scheme (resp. N -truncated simplicial scheme, for some $N \in \mathbb{N}$).

a) A k -augmentation $f_\bullet : X_\bullet \rightarrow X$ is called a *proper hypercovering* (resp. *N -truncated proper hypercovering*) of X if, for all $n \geq 0$ (resp. $0 \leq n \leq N$), the canonical morphism $X_n \rightarrow \mathrm{cosk}_{n-1}^X(\mathrm{sk}_{n-1}^X(X_\bullet))_n$ is proper and surjective.

b) A k -augmentation $f_\bullet : X_\bullet \rightarrow X$ is called an *étale hypercovering* (resp. *N -truncated étale hypercovering*) of X if, for all $n \geq 0$ (resp. $0 \leq n \leq N$), the canonical morphism $X_n \rightarrow \mathrm{cosk}_{n-1}^X(\mathrm{sk}_{n-1}^X(X_\bullet))_n$ is étale and surjective.

c) A k -augmentation $f_\bullet : X_\bullet \rightarrow X$ is called a *simplicial resolution* (resp. *N -truncated simplicial resolution*) of X if the following conditions hold:

(i) f_\bullet is a proper hypercovering (resp. N -truncated proper hypercovering) of X .

(ii) For all $n \geq 0$ (resp. $0 \leq n \leq N$), there exists a dense open subset $U_n \subset X$ such that the restriction of $\mathrm{sk}_n^X(X_\bullet)$ above U_n is a n -truncated étale hypercovering of U_n .

(iii) For all $n \geq 0$ (resp. $0 \leq n \leq N$), X_n is a smooth quasi-projective k -scheme.

Since k is perfect, de Jong's theorem [11, Th. 4.1 and Rem. 4.2] implies that, for any separated integral (*i.e.* reduced and irreducible) k -scheme of finite type X , there exists a quasi-projective, smooth and integral k -scheme X' , and a surjective, projective and generically étale morphism $X' \rightarrow X$. If X is separated and reduced, but not necessarily irreducible, one can apply de Jong's theorem to each irreducible component of X , and one gets in this way a quasi-projective smooth k -scheme X' , and a surjective projective morphism $X' \rightarrow X$ which is étale over a dense open subset of X . Replacing resolution of singularities by this result, one can then proceed as Deligne [12, (6.2.5)] to show that any separated reduced k -scheme of finite type has a simplicial resolution.

Remark. Let \mathcal{C} be the category of X -schemes which are separated of finite type over k , and \mathcal{D} the subcategory defined as follows:

- $\mathrm{Ob}(\mathcal{D}) = \mathrm{Ob}(\mathcal{C})$;
- If $Y, Z \in \mathrm{Ob}(\mathcal{D})$, $\mathrm{Hom}_{\mathcal{D}}(Y, Z)$ is the set of X -morphisms $f : Y \rightarrow Z$ which are proper, surjective, and such that there exists a dense open subset $U \subset X$ with the property that the restriction $f|_U : Y|_U \rightarrow Z|_U$ of f above U is étale.

Then \mathcal{D} satisfies the condition (HC) of [28, 5.1], and the simplicial resolutions of X are the \mathcal{D} -hypercoverings of X by smooth quasi-projective k -schemes, in the sense of [28, 5.1.1].

Proposition 3.2. *Let X be a reduced k -scheme of finite type, let $N \in \mathbb{N}$, and let $f_\bullet : X_\bullet \rightarrow X$ be a N -truncated proper hypercovering of X . Let $\mathcal{I} \subset \mathcal{O}_X$ be a coherent ideal, $\mathcal{I}_q = \mathcal{I}\mathcal{O}_{X_q}$, $Y = V(\mathcal{I})$, $U = X \setminus Y$, $U_q = f_q^{-1}(U)$. Assume that the restriction U_\bullet of X_\bullet above U is an N -truncated étale hypercovering of U . Then:*

(i) *The complex*

$$0 \rightarrow W\mathcal{I}_K \rightarrow f_{0*}(W\mathcal{I}_{0,K}) \rightarrow f_{1*}(W\mathcal{I}_{1,K}) \rightarrow \cdots \rightarrow f_{N*}(W\mathcal{I}_{N,K}),$$

where $W\mathcal{I}_K$ sits in degree -1 , is acyclic in degrees $\neq N$.

(ii) *The canonical morphisms*

$$(3.1) \quad H^q(X, W\mathcal{I}_K) \rightarrow H^q(X_\bullet, W\mathcal{I}_{\bullet,K})$$

are isomorphisms for all $q < N$.

Proof. For each $n \geq 1$ and each $r \geq 0$, we denote by $\mathcal{L}_n^{\bullet, r}$ the complex

$$(3.2) \quad 0 \rightarrow W_n(\mathcal{I}^r) \rightarrow f_{0*}(W_n(\mathcal{I}_0^r)) \rightarrow \cdots \rightarrow f_{N*}(W_n(\mathcal{I}_N^r)),$$

where $W_n(\mathcal{I}^r)$ sits in degree -1 , and we set

$$(3.3) \quad \mathcal{L}_n^\bullet = \bigoplus_{r \geq 0} \mathcal{L}_n^{\bullet, r},$$

$$(3.4) \quad \mathcal{H}_n^{q, r} = \mathcal{H}^q(\mathcal{L}_n^{\bullet, r}), \quad \mathcal{H}_n^q = \mathcal{H}^q(\mathcal{L}_n^\bullet) = \bigoplus_{r \geq 0} \mathcal{H}_n^{q, r}.$$

If we denote again by \mathcal{R} the graded \mathcal{O}_X -algebra $\bigoplus_{r \geq 0} \mathcal{I}^r$, it follows from [17, 3.3.1] that \mathcal{L}_1^\bullet is a complex of quasi-coherent graded \mathcal{R} -modules of finite type. Therefore, \mathcal{H}_1^q is a quasi-coherent graded \mathcal{R} -module of finite type for all q , and there exists an integer $d \geq 0$ such that the morphism

$$\mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{H}_1^{q, r} \rightarrow \mathcal{H}_1^{q, r+1}$$

is surjective for all $r \geq d$. Moreover, U_\bullet is an N -truncated étale hypercovering of U , and therefore it satisfies cohomological descent for quasi-coherent modules. It follows that $\mathcal{L}_1^\bullet|_U$ is acyclic in degrees $< N$. Then we can find an integer $m \geq 0$ such that $\mathcal{I}^m \mathcal{H}_1^{q, r} = 0$ for all $q < N$ and all $r \geq 0$. For $r \geq d$, this implies that the image of $\mathcal{H}_1^{q, r+m}$ in $\mathcal{H}_1^{q, r}$ is 0. Finally, if we set $b = d + m$, we obtain that the canonical morphism $\mathcal{H}_1^{q, r+b} \rightarrow \mathcal{H}_1^{q, r}$ is 0 for all $q < N$ and all $r \geq 0$.

We can now proceed as in the proof of Lemma 2.7 to prove that the morphism $\mathcal{H}_n^{q, r+b} \rightarrow \mathcal{H}_n^{q, r}$ is 0 for all $q \leq N$, all $n \geq 1$ and all $r \geq b$. As this last condition implies that $r \leq r + b \leq pr$, we can introduce for $n \geq 2$ the subcomplex $\mathcal{L}_n^{\bullet, r, r+b} \subset \mathcal{L}_n^{\bullet, r}$ defined by

$$0 \rightarrow W_n^{r, r+b}(\mathcal{I}) \rightarrow f_{0*}(W_n^{r, r+b}(\mathcal{I}_0)) \rightarrow \cdots \rightarrow f_{N*}(W_n^{r, r+b}(\mathcal{I}_N)),$$

and $\mathcal{H}_n^{q, r, r+b} = \mathcal{H}^q(\mathcal{L}_n^{\bullet, r, r+b})$. From Lemma 2.5, we deduce a commutative diagram of morphisms of complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{L}_{n-1}^{\bullet, r+b} & \xrightarrow{V} & \mathcal{L}_n^{\bullet, r+b} & \xrightarrow{R^{n-1}} & \mathcal{L}_1^{\bullet, r+b} \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{L}_{n-1}^{\bullet, r+b} & \xrightarrow{V} & \mathcal{L}_n^{\bullet, r, r+b}(I) & \xrightarrow{R^{n-1}} & \mathcal{L}_1^{\bullet, r} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{L}_{n-1}^{\bullet, r} & \xrightarrow{V} & \mathcal{L}_n^{\bullet, r} & \xrightarrow{R^{n-1}} & \mathcal{L}_1^{\bullet, r} \longrightarrow 0, \end{array}$$

where the rows are short exact sequences, because any section of a sheaf $f_{q*}(\mathcal{I}_q^r)$ can be lifted to a section of $f_{q*}(W_n(\mathcal{I}_q^r))$ or $f_{q*}(W_n^{r, r+b}(\mathcal{I}_q))$ by taking its Teichmüller representative. From the corresponding diagram

of cohomology exact sequences, we can extract for each q the commutative diagram

$$\begin{array}{ccccc}
& & \mathcal{H}_n^{q,r+b} & \xrightarrow{R^{n-1}} & \mathcal{H}_1^{q,r+b} \\
& & \downarrow & & \downarrow u_1 \\
\mathcal{H}_{n-1}^{q,r+b} & \xrightarrow{V} & \mathcal{H}_n^{q,r,r+b} & \xrightarrow{R^{n-1}} & \mathcal{H}_1^{q,r} \\
\downarrow u_{n-1} & & \downarrow & & \\
\mathcal{H}_{n-1}^{q,r} & \xrightarrow{V} & \mathcal{H}_n^{q,r} & & ,
\end{array}$$

where the morphisms u_1 , u_{n-1} and the composition of the two vertical morphisms are the canonical morphisms, and the middle row is exact. Since $u_1 = 0$ when $q < N$, Lemma 2.6 shows by induction that $u_n = 0$ for all $q < N$, all $n \geq 1$ and all $r \geq b$.

Setting $a = 2b$, we obtain that the morphisms $\mathcal{H}_n^{q,r+a} \rightarrow \mathcal{H}_n^{q,r}$ are 0 for all $q < N$, all $n \geq 1$ and all $r \geq 0$. This implies that, if c is chosen so that $p^c > a$, then F^c acts by 0 on the sheaves $\mathcal{H}_n^{q,r}$ for $r \geq 1$. Therefore, F^c acts by 0 on $R^i \varprojlim_n \mathcal{H}_n^{q,r}$ for all $i \geq 0$, all $q < N$ and all $r \geq 1$.

Let \mathcal{L}^\bullet be the complex

$$0 \rightarrow W\mathcal{I} \rightarrow f_{0*}(W\mathcal{I}_0) \rightarrow \cdots \rightarrow f_{N*}(W\mathcal{I}_N).$$

As the sheaves $\mathcal{L}_n^{q,1}$ have vanishing cohomology on affine open subsets of X , and the transition maps $\mathcal{L}_{n+1}^{q,1} \rightarrow \mathcal{L}_n^{q,1}$ are surjective, the projective systems $(\mathcal{L}_n^{q,1})_{n \geq 1}$ are \varprojlim_n -acyclic, and the canonical morphism

$$\mathcal{L}^\bullet \rightarrow \mathbb{R} \varprojlim_n \mathcal{L}_n^{\bullet,1}$$

is an isomorphism. This isomorphism provides a spectral sequence

$$E_2^{i,j} = R^i \varprojlim_n \mathcal{H}^j(\mathcal{L}_n^{\bullet,1}) \Rightarrow \mathcal{H}^{i+j}(\mathcal{L}^\bullet),$$

on which F acts. As F is nilpotent on the terms $E_2^{i,j}$ for $j < N$, it follows that F is also nilpotent on $\mathcal{H}^q(\mathcal{L}^\bullet)$ for $q < N$. But F becomes an isomorphism after tensorisation by K , so this implies assertion (i).

Thanks to Theorem 2.4, $R^i f_{q*}(W\mathcal{I}_q)_K = 0$ for all q and all $i \geq 1$. Therefore, the canonical morphism

$$f_{\bullet*}(W\mathcal{I}_\bullet)_K \rightarrow \mathbb{R} f_{\bullet*}(W\mathcal{I}_\bullet)_K$$

is an isomorphism. Then (ii) follows from (i). \square

4. WITT VECTOR COHOMOLOGY AND RIGID COHOMOLOGY

In this section, we first recall how to compute rigid cohomology in terms of de Rham complexes for the Zariski topology. Using this description, we construct the canonical morphism from rigid cohomology with compact supports to Witt vector cohomology with compact supports.

4.1. We begin by recalling the general construction of tubes [3], which is sufficient to define the rigid cohomology groups of a proper k -scheme.

Let X be a separated k -scheme of finite type, and let $X \hookrightarrow \mathbb{P}$ be a closed immersion into a smooth formal W -scheme. The formal scheme \mathbb{P} has a generic fibre \mathbb{P}_K , which is a rigid analytic space, endowed with a continuous morphism $\mathrm{sp} : \mathbb{P}_K \rightarrow \mathbb{P}$, the specialization morphism, such that $\mathrm{sp}^{-1}(\mathbb{U}) = \mathbb{U}_K$ for any open subset $\mathbb{U} \subset \mathbb{P}$, and $\mathrm{sp}_* \mathcal{O}_{\mathbb{P}_K} = \mathcal{O}_{\mathbb{P},K}$. When \mathbb{P} is an affine formal scheme $\mathrm{Spf}A$, where A is a W -algebra which is topologically of finite type, its generic fibre is the affinoid space $\mathrm{Spm}A_K$, defined by the Tate algebra A_K .

If $\mathbb{P} = \mathrm{Spf}A$, and if $f_1, \dots, f_r \in A$ is a family of generators of the ideal \mathcal{J} of X in \mathbb{P} , the tube $]X[_{\mathbb{P}}$ of X in \mathbb{P}_K is the admissible open subset defined by

$$]X[_{\mathbb{P}} = \{x \in \mathbb{P}_K \mid \forall i, |f_i(x)| < 1\},$$

where, for a point $x \in \mathbb{P}_K$ corresponding to a maximal ideal $\mathfrak{m} \subset A_K$, $|f_i(x)|$ is the absolute value of the class of f_i in the residue field $K(x) = A_K/\mathfrak{m}$. When \mathbb{P} is not affine, one can choose a covering of \mathbb{P} by open affine subsets \mathbb{U}_i , and the generic fibres $\mathbb{U}_{i,K}$ of the \mathbb{U}_i provide an admissible covering of \mathbb{P}_K . Then the tube $]X[_{\mathbb{P}}$ can be defined by gluing the tubes $]X \cap \mathbb{U}_i[_{\mathbb{U}_i} \subset \mathbb{U}_{i,K}$.

4.2. We introduce now the $\mathcal{O}_{\mathbb{P}}$ -algebra $\mathcal{A}_{X,\mathbb{P}}$ of analytic functions on $]X[_{\mathbb{P}}$, the de Rham cohomology of which defines rigid cohomology.

The specialization morphism $\mathrm{sp} : \mathbb{P}_K \rightarrow \mathbb{P}$ maps $]X[_{\mathbb{P}}$ to X . Thus, one can define a sheaf of $\mathcal{O}_{\mathbb{P},K}$ -algebras supported on X by setting

$$\mathcal{A}_{X,\mathbb{P}} = \mathrm{sp}_* \mathcal{O}_{]X[_{\mathbb{P}}}.$$

The differentiation of analytic functions on $]X[_{\mathbb{P}}$ endows $\mathcal{A}_{X,\mathbb{P}}$ with a canonical integrable connection, allowing to define the de Rham complex $\mathcal{A}_{X,\mathbb{P}} \otimes \Omega_{\mathbb{P}}^{\bullet}$. For any affine open subset $\mathbb{U} \subset \mathbb{P}$ and any j , we have $\Gamma(\mathbb{U}_K, \Omega_{\mathbb{P}_K}^j) \cong \Gamma(\mathbb{U}, \Omega_{\mathbb{P}}^j) \otimes K$. It follows that there is a canonical isomorphism of complexes

$$\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \cong \mathrm{sp}_*(\Omega_{]X[_{\mathbb{P}}}^{\bullet}).$$

Moreover, the above description of $]X[_{\mathbb{P}}$ in the affine case shows that the inverse image of an affine open subset $U \subset \mathbb{P}$ is quasi-Stein in the sense of Kiehl [22, Definition 2.3]. Therefore, Kiehl's vanishing theorem for coherent analytic sheaves implies that $R^i \mathrm{sp}_*(\Omega_{]X[_{\mathbb{P}}}^j) = 0$ for all j and all $i \geq 1$, and we obtain

$$\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \cong \mathbb{R}\mathrm{sp}_*(\Omega_{]X[_{\mathbb{P}}}^{\bullet})$$

in the derived category $D^b(X, K)$.

If $X \hookrightarrow \mathbb{P}$ and $X \hookrightarrow \mathbb{P}'$ are two closed immersions of X into smooth formal W -schemes, there exists in $D^b(X, K)$ a canonical isomorphism [4, 1.5]

$$(4.1) \quad \mathcal{A}_{X,\mathbb{P}} \otimes \Omega_{\mathbb{P}}^{\bullet} \cong \mathcal{A}_{X,\mathbb{P}'} \otimes \Omega_{\mathbb{P}'}^{\bullet}.$$

Thus, up to canonical isomorphism, the complex $\mathcal{A}_{X,\mathbb{P}} \otimes \Omega_{\mathbb{P}}^{\bullet}$ does not depend in $D^b(X, K)$ on the choice of the embedding $X \hookrightarrow \mathbb{P}$. In particular, the de Rham cohomology of $]X[_{\mathbb{P}}$ does not depend on this choice. When X is a proper k -scheme and can be embedded in a smooth formal scheme \mathbb{P} as above, its rigid cohomology is defined by

$$(4.2) \quad \mathbb{R}\Gamma_{\mathrm{rig}}(X/K) = \mathbb{R}\Gamma(]X[_{\mathbb{P}}, \Omega_{]X[_{\mathbb{P}}}^{\bullet}) \cong \mathbb{R}\Gamma(X, \mathcal{A}_{X,\mathbb{P}} \otimes \Omega_{\mathbb{P}}^{\bullet}).$$

If such an embedding does not exist, one can choose a covering of X by affine open subsets X_i , and, for each i , a closed immersion of X_i in a smooth affine formal scheme \mathbb{P}_i over W . Using the diagonal immersions for finite intersections $X_{i_0, \dots, i_r} = X_{i_0} \cap \dots \cap X_{i_r}$, and the corresponding algebras $\mathcal{A}_{X_{i_0, \dots, i_r}, \mathbb{P}_{i_0} \times \dots \times \mathbb{P}_{i_r}}$, one can build a Čech-de Rham double complex. When X is proper, the cohomology of the associated total complex defines rigid cohomology.

4.3. The previous constructions can be extended as follows to define rigid cohomology with compact supports for separated k -schemes of finite type.

Let U be such a scheme, $U \hookrightarrow X$ an open immersion of U in a proper k -scheme X , $Y = X \setminus U$. Assume that there exists a closed immersion $X \hookrightarrow \mathbb{P}$ of X into a smooth formal scheme \mathbb{P} , and denote by $u :]Y[_{\mathbb{P}} \hookrightarrow]X[_{\mathbb{P}}$ the inclusion of the tube of Y into the tube of X . Then $]Y[_{\mathbb{P}}$ is an admissible open subset of $]X[_{\mathbb{P}}$, and, by construction [2], the rigid cohomology with compact supports of U is defined by

$$\mathbb{R}\Gamma_{\mathrm{rig},c}(U/K) = \mathbb{R}\Gamma(]X[_{\mathbb{P}}, (\Omega_{]X[_{\mathbb{P}}}^{\bullet} \rightarrow u_*(\Omega_{]Y[_{\mathbb{P}}}^{\bullet}))_t),$$

where the subscript t denotes the total complex associated to a double complex. Using the algebras $\mathcal{A}_{X,\mathbb{P}}$ and $\mathcal{A}_{Y,\mathbb{P}}$ defined in 4.2, we can

rephrase this definition as

$$(4.3) \quad \mathbb{R}\Gamma_{\text{rig,c}}(U/K) = \mathbb{R}\Gamma(X, (\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow \mathcal{A}_{Y,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet})_t).$$

This cohomology only depends on U , and has the usual properties of cohomology with compact supports.

If there is no embedding of a compactification X of U in a smooth formal W -scheme, the rigid cohomology with compact supports of U can still be defined using coverings by affine open subsets and Čech complexes as in 4.2.

Proposition 4.4. *Let \mathbb{P} be a smooth and separated formal scheme over W , with special fibre P , and let X be a closed subscheme of P .*

(i) *The datum of a σ -semilinear lifting $F : \mathbb{P} \rightarrow \mathbb{P}$ of the absolute Frobenius endomorphism of P defines a ring homomorphism*

$$(4.4) \quad \mathcal{A}_{X,\mathbb{P}} \rightarrow W\mathcal{O}_{X,K},$$

functorial in (X, \mathbb{P}, F) .

(ii) *Without assumption on the existence of F , there exists in $D^b(X, K)$ a morphism*

$$(4.5) \quad a_{X,\mathbb{P}} : \mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow W\mathcal{O}_{X,K},$$

functorial in (X, \mathbb{P}) , equal to the composed morphism

$$\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow \mathcal{A}_{X,\mathbb{P}} \xrightarrow{(4.4)} W\mathcal{O}_{X,K}$$

whenever there exists a lifting of the Frobenius endomorphism on \mathbb{P} , and compatible with the canonical isomorphism (4.1)

$$\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \cong \mathcal{A}_{X,\mathbb{P}'} \otimes_{\mathcal{O}_{\mathbb{P}'}} \Omega_{\mathbb{P}'}^{\bullet}$$

for two closed immersions of X into smooth and separated formal schemes \mathbb{P} and \mathbb{P}' .

(iii) *Let U be an open subset of X , $Y = X \setminus U$, and $\mathcal{I} \subset \mathcal{O}_X$ a coherent ideal such that $V(\mathcal{I}) = Y$. Without assumption on the existence of F , there exists in $D^b(X, K)$ a morphism*

$$(4.6) \quad a_{U,X,\mathbb{P}} : (\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow \mathcal{A}_{Y,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet})_t \rightarrow W\mathcal{I}_K,$$

functorial with respect to morphisms $(U', X', \mathbb{P}') \rightarrow (U, X, \mathbb{P})$ such that $U' \rightarrow U$ is proper, equal to (4.5) when $U = X$, and compatible with the canonical isomorphisms

$$(\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow \mathcal{A}_{Y,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet})_t \cong (\mathcal{A}_{X,\mathbb{P}'} \otimes_{\mathcal{O}_{\mathbb{P}'}} \Omega_{\mathbb{P}'}^{\bullet} \rightarrow \mathcal{A}_{Y,\mathbb{P}'} \otimes_{\mathcal{O}_{\mathbb{P}'}} \Omega_{\mathbb{P}'}^{\bullet})_t$$

for two closed immersions of X into smooth formal schemes \mathbb{P} and \mathbb{P}' .

Proof. Let $\mathcal{J} \subset \mathcal{O}_{\mathbb{P}}$ be the ideal defining X , $\mathcal{P}(\mathcal{J})$ the divided power envelope of \mathcal{J} , with compatibility with the natural divided powers of p , and $\widehat{\mathcal{P}}(\mathcal{J})$ its p -adic completion. We recall first from [4, 1.9] that there exists a functorial homomorphism of $\mathcal{O}_{\mathbb{P}}$ -algebras

$$(4.7) \quad \mathcal{A}_{X, \mathbb{P}} \rightarrow \widehat{\mathcal{P}}(\mathcal{J})_K.$$

Let P_n be the reduction of $\mathbb{P} \bmod p^n$, and let $\mathcal{J}_n = \mathcal{J}\mathcal{O}_{P_n}$ be the ideal of X in P_n . Thanks to the compatibility condition with the divided powers of p , there is a canonical isomorphism

$$\widehat{\mathcal{P}}(\mathcal{J})/p^n \widehat{\mathcal{P}}(\mathcal{J}) \xrightarrow{\sim} \mathcal{P}(\mathcal{J}_n),$$

where $\mathcal{P}(\mathcal{J}_n)$ is the divided power envelope of \mathcal{J}_n with the same compatibility condition, and we obtain

$$\widehat{\mathcal{P}}(\mathcal{J}) \xrightarrow{\sim} \varprojlim_n \mathcal{P}(\mathcal{J}_n).$$

Thus, it suffices to define a compatible family of functorial ring homomorphisms

$$(4.8) \quad \mathcal{P}(\mathcal{J}_n) \rightarrow W_n \mathcal{O}_X$$

to obtain a morphism (4.4).

Let us assume that \mathbb{P} is endowed with a lifting F of the Frobenius morphism. As $\mathcal{O}_{\mathbb{P}}$ is p -torsion free, the homomorphism $F : \mathcal{O}_{\mathbb{P}} \rightarrow \mathcal{O}_{\mathbb{P}}$ defines a section $s_F : \mathcal{O}_{\mathbb{P}} \rightarrow W\mathcal{O}_{\mathbb{P}}$ of the reduction homomorphism $W\mathcal{O}_{\mathbb{P}} \rightarrow \mathcal{O}_{\mathbb{P}}$, characterized by the fact that $w_i(s_F(x)) = F^i(x)$ for any $x \in \mathcal{O}_{\mathbb{P}}$ and any ghost component w_i [20, 0 1.3]. Composing s_F with the homomorphisms $W\mathcal{O}_{\mathbb{P}} \rightarrow W\mathcal{O}_P \rightarrow W_n \mathcal{O}_P$ and factorizing, we obtain for all $n \geq 1$ a homomorphism

$$\mathcal{O}_{P_n} \rightarrow W_n \mathcal{O}_P \rightarrow W_n \mathcal{O}_X,$$

which maps \mathcal{J}_n to $VW_{n-1} \mathcal{O}_X \subset W_n \mathcal{O}_X$. The ideal $VW_{n-1} \mathcal{O}_X$ has a natural structure of divided power ideal (compatible with the divided powers of p), defined by $(Vx)^{[i]} = (p^{i-1}/i!)V(x^i)$ for all $i \geq 1$. Therefore, this homomorphism factors through a homomorphism $\mathcal{P}(\mathcal{J}_n) \rightarrow W_n \mathcal{O}_X$. This provides the compatible family of homomorphisms defining (4.4), and it is clear that the homomorphism obtained in this way is functorial in (X, \mathbb{P}, F) .

By composition with the augmentation morphism $\mathcal{A}_{X, \mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow \mathcal{A}_{X, \mathbb{P}}$, we obtain a morphism of complexes

$$a_{X, \mathbb{P}, F} : \mathcal{A}_{X, \mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow W\mathcal{O}_{X, K},$$

which is still functorial in (X, \mathbb{P}, F) . If $X \hookrightarrow \mathbb{P}$ and $X \hookrightarrow \mathbb{P}'$ are two closed immersions in smooth formal schemes endowed with liftings

of Frobenius F and F' , we can endow $\mathbb{P}'' = \mathbb{P} \times_W \mathbb{P}'$ with $F \times F'$, and embed X diagonally into \mathbb{P}'' . Applying the functoriality of this construction to the two projections, we obtain a commutative diagram

$$(4.9) \quad \begin{array}{ccccc} \mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} & \longrightarrow & \mathcal{A}_{X,\mathbb{P}''} \otimes_{\mathcal{O}_{\mathbb{P}''}} \Omega_{\mathbb{P}''}^{\bullet} & \longleftarrow & \mathcal{A}_{X,\mathbb{P}'} \otimes_{\mathcal{O}_{\mathbb{P}'}} \Omega_{\mathbb{P}'}^{\bullet} \\ \downarrow a_{X,\mathbb{P},F} & & \downarrow a_{X,\mathbb{P}'',F''} & & \downarrow a_{X,\mathbb{P}',F'} \\ W\mathcal{O}_{X,K} & \xlongequal{\quad} & W\mathcal{O}_{X,K} & \xlongequal{\quad} & W\mathcal{O}_{X,K} \end{array}$$

in which the morphisms of the top row are quasi-isomorphisms. In $D^b(X, K)$, the composed isomorphism $\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \cong \mathcal{A}_{X,\mathbb{P}'} \otimes_{\mathcal{O}_{\mathbb{P}'}} \Omega_{\mathbb{P}'}^{\bullet}$ is the canonical isomorphism (4.1) [4, 1.5], which shows the compatibility asserted in (ii). In the particular case where $\mathbb{P}' = \mathbb{P}$, this composed isomorphism is the identity, and we obtain that, as a morphism of $D^b(X, K)$, the morphism $a_{X,\mathbb{P},F}$ does not depend on the choice of F . A similar argument shows that, in $D^b(X, K)$, it is functorial with respect to morphisms $(X', \mathbb{P}') \rightarrow (X, \mathbb{P})$ without compatibility with Frobenius liftings. Thus, if we define $a_{X,\mathbb{P}}$ to be the image in $D^b(X, K)$ of $a_{X,\mathbb{P},F}$, assertion (ii) is true when there exists a lifting of Frobenius on \mathbb{P} .

In the general case, we can choose an affine covering \mathbb{P}_i of \mathbb{P} , and a lifting of Frobenius F_i on each \mathbb{P}_i . Then, if $X_{i_0, \dots, i_n} = X \cap \mathbb{P}_{i_0} \cap \dots \cap \mathbb{P}_{i_n}$ and j_{i_0, \dots, i_n} denotes its inclusion in X , the complex $\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet}$ is quasi-isomorphic to the total complex associated to the double complex

$$(4.10) \quad \begin{aligned} & \prod_i j_{i*}(\mathcal{A}_{X_i, \mathbb{P}_i} \otimes \Omega_{\mathbb{P}_i}^{\bullet}) \rightarrow \dots \\ & \rightarrow \prod_{i_0, \dots, i_n} j_{i_0, \dots, i_n*}(\mathcal{A}_{X_{i_0, \dots, i_n}, \mathbb{P}_{i_0} \times \dots \times \mathbb{P}_{i_n}} \otimes \Omega_{\mathbb{P}_{i_0} \times \dots \times \mathbb{P}_{i_n}}^{\bullet}) \rightarrow \dots, \end{aligned}$$

while $W\mathcal{O}_{X,K}$ is quasi-isomorphic to the Čech resolution

$$(4.11) \quad \prod_i j_{i*}(W\mathcal{O}_{X_i, K}) \rightarrow \dots \rightarrow \prod_{i_0, \dots, i_n} j_{i_0, \dots, i_n*}(W\mathcal{O}_{X_{i_0, \dots, i_n}, K}) \rightarrow \dots.$$

Then we can define $a_{X,\mathbb{P}}$ as the image in $D^b(X, K)$ of the morphism defined by the collection of all $a_{X_{i_0, \dots, i_n}, \mathbb{P}_{i_0} \times \dots \times \mathbb{P}_{i_n}, F_{i_0} \times \dots \times F_{i_n}}$, and assertion (ii) is verified as above.

Let U be an open subset of X , and assume again that there exists a lifting F of the Frobenius endomorphism on \mathbb{P} . Then the previous constructions can be applied both to X and to $Y = X \setminus U$. By functoriality, they provide a morphism of complexes

$$(\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet} \rightarrow \mathcal{A}_{Y,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^{\bullet})_t \rightarrow (W\mathcal{O}_{X,K} \rightarrow W\mathcal{O}_{Y,K}).$$

As the length 1 complex $W\mathcal{O}_{X,K} \rightarrow W\mathcal{O}_{Y,K}$ is a resolution of $W\mathcal{I}_K$, we can define $a_{U,X,\mathbb{P}}$ as the image of this morphism in $D^b(X, K)$. If $Y = \emptyset$, then we simply obtain (4.5). To check the functoriality, we may assume that U' is dense in X' , because $W\mathcal{I}'_K$ does not change if we replace X' by the closure of U' in X' , by 2.4, and similarly $(\mathcal{A}_{X,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^\bullet \rightarrow \mathcal{A}_{Y,\mathbb{P}} \otimes_{\mathcal{O}_{\mathbb{P}}} \Omega_{\mathbb{P}}^\bullet)_t$ does not change up to canonical isomorphism in $D^b(X, K)$, by the basic properties of rigid cohomology. Then the properness of $U' \rightarrow U$ allows to assume that $Y' = X' \setminus U'$ is defined by $\mathcal{I}' = \mathcal{I}\mathcal{O}_{X'}$, and the functoriality is clear. Finally, the compatibility with the canonical isomorphism for two embeddings of X into smooth formal schemes results from the same assertion for the morphisms (4.5) relative to X and Y .

When F cannot be lifted to \mathbb{P} , one can proceed with Čech coverings as in (ii) to define (4.6), and the same properties hold. \square

Remark. If X is quasi-projective, one can always find a closed immersion of X into a smooth formal scheme \mathbb{P} endowed with a lifting of Frobenius F , since it suffices to choose for \mathbb{P} an open subscheme of a projective space, endowed with the endomorphism induced by some lifting of the Frobenius endomorphism of the projective space.

Theorem 4.5. *Let U be a separated k -scheme of finite type. In $D^b(K)$, there exists a functorial morphism*

$$(4.12) \quad a_U : \mathbb{R}\Gamma_{\text{rig,c}}(U/K) \rightarrow \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K}),$$

equal, for a proper and smooth k -scheme X , to the canonical morphism

$$(4.13) \quad \mathbb{R}\Gamma_{\text{rig}}(X/K) \xrightarrow{\sim} \mathbb{R}\Gamma_{\text{crys}}(X/W)_K \xrightarrow{\sim} \mathbb{R}\Gamma(X, W\Omega_X^\bullet)_K$$

$$\downarrow$$

$$\mathbb{R}\Gamma(X, W\mathcal{O}_X)_K.$$

In (4.13), the first isomorphism is the comparison isomorphism between rigid and crystalline cohomologies [4, 1.9], the second one is the comparison isomorphism between crystalline and de Rham-Witt cohomologies [20, II, (1.3.2)], and the third map is defined by the augmentation morphism of the de Rham-Witt complex.

Proof. Let $U \hookrightarrow X$ be an open immersion of U in a proper k -scheme. Let us assume first that there exists a closed immersion $X \hookrightarrow \mathbb{P}$ of X into a smooth formal scheme \mathbb{P} . To define a_U , we apply the functor $\mathbb{R}\Gamma(X, -)$ to the morphism $a_{U,X,\mathbb{P}}$ defined in (4.6). Because of the compatibility property of 4.4 (iii), we obtain in this way a morphism of $D^b(K)$ which does not depend, up to canonical isomorphism, on the

choice of \mathbb{P} . Using the functoriality of $a_{U,X,\mathbb{P}}$, it is easy to check that it does not depend either on the choice of the compactification X of U , and that it is functorial with respect to U .

When $U = X$ is a proper and smooth k -scheme, and X is embeddable in a smooth formal scheme \mathbb{P} with a lifting F of the Frobenius endomorphism, the homomorphisms (4.8) used to construct $a_{X,\mathbb{P},F}$ are the homomorphisms used in [20, II, (1.1.5)] to construct the morphism of complexes $\widehat{\mathcal{P}}(\mathcal{J}) \otimes \Omega_{\mathbb{P}}^{\bullet} \rightarrow W\Omega_X^{\bullet}$ which defines the isomorphism between crystalline and de Rham-Witt cohomologies. Therefore, the comparison isomorphisms which appear in (4.13) sit in a commutative diagram

$$\begin{array}{ccccc} \mathbb{R}\Gamma(X, \mathcal{A}_{X,\mathbb{P}} \otimes \Omega_{\mathbb{P}}^{\bullet}) & \xrightarrow{\sim} & \mathbb{R}\Gamma(X, \widehat{\mathcal{P}}(\mathcal{J}) \otimes \Omega_{\mathbb{P},K}^{\bullet}) & \xrightarrow{\sim} & \mathbb{R}\Gamma(X, W\Omega_{X,K}^{\bullet}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{R}\Gamma(X, \mathcal{A}_{X,\mathbb{P}}) & \longrightarrow & \mathbb{R}\Gamma(X, \widehat{\mathcal{P}}(\mathcal{J})_K) & \longrightarrow & \mathbb{R}\Gamma(X, W\mathcal{O}_{X,K}), \end{array}$$

in which the vertical maps are induced by the augmentation morphisms of the complexes appearing in the upper row. The equality of the morphisms (4.12) and (4.13) follows immediately.

In the general case where there is no smooth embedding \mathbb{P} of X , or no Frobenius lifting on \mathbb{P} , one can again use a covering of X by affine open subsets X_i , and closed immersions $X_i \hookrightarrow \mathbb{P}_i$ in smooth formal schemes \mathbb{P}_i endowed with liftings of Frobenius, to construct Čech complexes as in (4.10) and (4.11). One can then use on each intersection of the covering the corresponding morphism (4.6) and define in this way a morphism between the two Čech complexes. Applying $\mathbb{R}\Gamma(X, -)$, one gets in $D^b(K)$ the morphism a_U . In the proper and smooth case, the fact that each map in the above diagram can be defined through Čech complexes shows that a_U is still equal to (4.13). \square

The following verifications are left as an exercise for the reader.

Proposition 4.6. *The morphism a_U defined by Theorem 4.5 satisfies the following compatibility properties.*

(i) *The diagram*

$$(4.14) \quad \begin{array}{ccc} \mathbb{R}\Gamma_{\text{rig,c}}(U/K) & \xrightarrow{\sim} & \mathbb{R}\Gamma_{\text{rig,c}}(U_{\text{red}}/K) \\ \downarrow a_U & & \downarrow a_{U_{\text{red}}} \\ \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U,K}) & \xrightarrow{\sim} & \mathbb{R}\Gamma_c(U, W\mathcal{O}_{U_{\text{red}},K}) \end{array}$$

is commutative.

(ii) If $V \subset U$ is an open subset, and $T = U \setminus V$, the morphisms a_V , a_U and a_T define a morphism of exact triangles

$$(4.15) \quad \begin{array}{ccccccc} \mathbb{R}\Gamma_{\text{rig},c}(V/K) & \longrightarrow & \mathbb{R}\Gamma_{\text{rig},c}(U/K) & \longrightarrow & \mathbb{R}\Gamma_{\text{rig},c}(T/K) & \xrightarrow{+1} & \longrightarrow \\ \downarrow a_V & & \downarrow a_U & & \downarrow a_T & & \\ R\Gamma_c(V, W\mathcal{O}_{V,K}) & \longrightarrow & R\Gamma_c(U, W\mathcal{O}_{U,K}) & \longrightarrow & R\Gamma_c(T, W\mathcal{O}_{T,K}) & \xrightarrow{+1} & \longrightarrow \end{array} .$$

5. PROOF OF THE MAIN THEOREM

We prove here that the morphism a_U constructed in Theorem 4.5 yields an identification of the slope < 1 subspace of rigid cohomology with compact supports with Witt vector cohomology with compact supports, thus completing the proof of Theorem 1.1.

The next lemma will allow us use descent techniques to study the morphism a_U . We follow here the method of Chiarellotto and Tsuzuki to construct embeddings of a simplicial scheme in a smooth simplicial formal scheme (cf. [10, 11.2], [28, 7.3]).

Lemma 5.1. *Let X be a separated reduced k -scheme of finite type, and let $X \hookrightarrow \mathbb{P}$ be a closed immersion into a smooth formal W -scheme endowed with a Frobenius lifting F . Let $N \in \mathbb{N}$ be a fixed integer. There exists a proper hypercovering X_\bullet of X , a \mathbb{P} -augmented simplicial formal W -scheme \mathbb{P}_\bullet , endowed with a σ -semilinear endomorphism F_\bullet lifting the absolute Frobenius endomorphism of its special fibre, and a morphism of simplicial schemes $X_\bullet \rightarrow \mathbb{P}_\bullet$ above $X \hookrightarrow \mathbb{P}$, such that the following conditions are satisfied:*

(i) *For all n , the morphism $X_n \rightarrow \mathbb{P}_n$ is a closed immersion, and the projection $\mathbb{P}_n \rightarrow \mathbb{P}$ commutes with the Frobenius liftings.*

(ii) *The N -truncated simplicial scheme $\text{sk}_N^X(X_\bullet)$ is a N -truncated simplicial resolution of X .*

(iii) *The canonical morphisms $X_\bullet \rightarrow \text{cosk}_N^X(\text{sk}_N^X(X_\bullet))$ and $\mathbb{P}_\bullet \rightarrow \text{cosk}_N^{\mathbb{P}}(\text{sk}_N^{\mathbb{P}}(\mathbb{P}_\bullet))$ are isomorphisms.*

(iv) *For all n , \mathbb{P}_n is smooth over \mathbb{P} , and the canonical morphism $\mathbb{P}_n \rightarrow \text{cosk}_{n-1}^{\mathbb{P}}(\text{sk}_{n-1}^{\mathbb{P}}(\mathbb{P}_\bullet))_n$ is smooth.*

Proof. Using de Jong's theorem, we can find an N -truncated simplicial resolution of X (cf. 3.1). We define X_\bullet as the N -coskeleton over X of this N -truncated resolution. Thus X_\bullet is a proper hypercovering of X , $\text{sk}_N^X(X_\bullet)$ is our initial N -truncated simplicial resolution of X , and the morphism $X_\bullet \rightarrow \text{cosk}_N^X(\text{sk}_N^X(X_\bullet))$ is an isomorphism.

For $n \in \mathbb{N}$, let $[n]$ denote the ordered set $\{0, \dots, n\}$. Let Δ be the category which has the sets $[n]$ as objects, and the set of non decreasing maps $[m] \rightarrow [n]$ as set of morphisms from $[m]$ to $[n]$. Since the X_m are quasi projective over k , we can choose for each $m \leq N$ a closed immersion $i_m : X_m \hookrightarrow \mathbb{W}_m$, where \mathbb{W}_m is a smooth formal W -scheme endowed with a lifting of Frobenius F_m . From \mathbb{W}_m , we construct a simplicial complex of smooth formal W -schemes $\Gamma_n(\mathbb{W}_m)$ by setting for all $n \geq 0$

$$(5.1) \quad \Gamma_n(\mathbb{W}_m) = \prod_{\gamma: [m] \rightarrow [n]} \mathbb{W}_{m, \gamma},$$

where the product is taken over all morphisms $\gamma : [m] \rightarrow [n]$ in Δ , and $\mathbb{W}_{m, \gamma} = \mathbb{W}_m$ for all γ . If $\eta : [n'] \rightarrow [n]$ is a morphism in Δ , the corresponding morphism $\Gamma_n(\mathbb{W}_m) \rightarrow \Gamma_{n'}(\mathbb{W}_m)$ is such that, for any $\gamma' : [m] \rightarrow [n']$ in Δ , its composition with the projection $\Gamma_{n'}(\mathbb{W}_m) \rightarrow \mathbb{W}_{m, \gamma'} = \mathbb{W}_m$ is the projection of $\Gamma_n(\mathbb{W}_m)$ on $\mathbb{W}_{m, \eta \circ \gamma'} = \mathbb{W}_m$. We can then define a Frobenius lifting on the simplicial formal scheme $\Gamma_\bullet(\mathbb{W}_m)$ as being the product morphism defined by F_m on each $\Gamma_n(\mathbb{W}_m)$; we still denote it by F_m .

The immersion i_m defines a morphism of simplicial schemes $X_\bullet \rightarrow \Gamma_\bullet(\mathbb{W}_m)$ as follows: for each $n \geq 0$, the composition of $X_n \rightarrow \Gamma_n(\mathbb{W}_m)$ with the projection of index γ is the morphism $X_n \rightarrow X_m \hookrightarrow \mathbb{W}_m$, where the first morphism is the morphism of X_\bullet defined by γ . We can now define a simplicial formal W -scheme \mathbb{P}_\bullet augmented towards \mathbb{P} by setting

$$(5.2) \quad \mathbb{P}_\bullet = \mathbb{P} \times_{\mathrm{Spf}(W)} \prod_{0 \leq m \leq N} \Gamma_\bullet(\mathbb{W}_m).$$

We define a morphism of simplicial schemes $X_\bullet \rightarrow \mathbb{P}_\bullet$ using the composed morphism $X_\bullet \rightarrow X \rightarrow \mathbb{P}$ and the family of morphisms $X_\bullet \rightarrow \Gamma_\bullet(\mathbb{W}_m)$ defined above. By construction, this morphism is compatible with the immersion $X \hookrightarrow \mathbb{P}$ via the augmentation morphisms. If we endow \mathbb{P}_\bullet with the Frobenius lifting $F \times \prod_m F_m$, then the augmentation $\mathbb{P}_\bullet \rightarrow \mathbb{P}$ commutes with the Frobenius liftings.

To check the remaining properties, we observe first that $\mathbb{W}_\bullet \xrightarrow{\sim} \mathrm{cosk}_N^{\mathbb{P}}(\mathrm{sk}_N^{\mathbb{P}}(\mathbb{W}_\bullet))$ thanks to [10, 11.2.5]. For each $m \leq N$, the morphism $X_m \rightarrow \Gamma_m(\mathbb{W}_m)$ is a closed immersion since the chosen immersion $X_m \hookrightarrow \mathbb{W}_m$ is one of the factors. Therefore $X_n \rightarrow \mathbb{P}_n$ is a closed immersion for all $n \leq N$. Then, thanks to the previous property, it follows from the construction of coskeletons that $X_n \rightarrow \mathbb{P}_n$ is a closed

immersion for all n . Finally, each \mathbb{P}_n is smooth over \mathbb{P} by construction, and the fact that \mathbb{P}_n is smooth over $(\text{cosk}_{n-1}^{\mathbb{P}}(\text{sk}_{n-1}^{\mathbb{P}}(\mathbb{P}\bullet)))_n$ for all n follows from [28, 7.3.3]. \square

5.2. Proof of Theorem 1.1. We first recall the definition of the subspaces with prescribed slopes. Let \bar{k} be an algebraic closure of k , and let $\bar{K}_0 = \text{Frac}(W(\bar{k}))$. If E is a finite dimensional K -vector space endowed with a σ -semilinear automorphism F , let $\bar{E} = \bar{K}_0 \otimes_K E$, and $\bar{F} = \sigma_{\bar{K}_0} \otimes F$. The Dieudonné-Manin theorem provides for each $\lambda \in \mathbb{Q}$ a maximal \bar{F} -stable subspace \bar{E}^λ on which \bar{F} has purely slope λ , and asserts that $\bigoplus_{\lambda} \bar{E}^\lambda \cong E$. As these subspaces are invariant under the action of $\text{Gal}(\bar{k}/k)$, there exists a unique F -stable subspace $E^\lambda \subset E$ such that $\bar{K}_0 \otimes_K E^\lambda \xrightarrow{\sim} \bar{E}^\lambda$, and $\bigoplus_{\lambda} E^\lambda \cong E$. The subspace E^λ is the *slope λ subspace* of E , and, for any $\rho \in \mathbb{R}$, we define the *slope $< \rho$ subspace* of E by

$$E^{<\rho} = \bigoplus_{\lambda < \rho} E^\lambda.$$

These spaces are functorial with respect to (E, F) , and define exact functors.

Let X be a separated k -scheme of finite type. The morphism (4.12) induces canonical homomorphisms

$$a_X^q : H_{\text{rig,c}}^q(X/K) \rightarrow H_c^q(X, W\mathcal{O}_{X,K})$$

between the cohomology spaces. By functoriality, a_X^q commutes to the Frobenius actions on both sides. The spaces $H_{\text{rig,c}}^q(X/K)$ are finite dimensional [4, 3.9 (i)], and their Frobenius endomorphism is an automorphism [16, 2.1]. Therefore they have a slope decomposition, and a_X^q induces a homomorphism

$$b_X^q : H_{\text{rig,c}}^q(X/K)^{<1} \rightarrow H_c^q(X, W\mathcal{O}_{X,K})$$

which is the homomorphism (1.3) of Theorem 1.1.

To prove that b_X^q is an isomorphism, we first observe that this is true when X is proper and smooth over k . Indeed, a_X^q is then induced by the morphism (4.13), and the isomorphisms which enter in the definition of (4.13) are compatible with Frobenius, hence induce isomorphisms on the slope < 1 subspaces of the cohomology spaces. The theorem is then a consequence of the slope decomposition for de Rham-Witt cohomology [20, II, (3.5.2)].

In the general case, we prove Theorem 1.1 by induction on $\dim(X)$. The diagram (4.14) allows us to assume that X is reduced. In particular, the theorem holds for $\dim(X) = 0$ thanks to the previous remark.

Let us assume that all b_X^q are isomorphisms when $\dim(X) < d$. If $U \subset X$ is an open subset of dimension d of a proper and smooth k -scheme X , its closure \bar{U} is proper and smooth over k , and $\dim(\bar{U} \setminus U) < d$. As all $b_{\bar{U}}^q$ are isomorphisms, the induction hypothesis and the commutativity of (4.15) imply that all b_U^q are isomorphisms. Thus the theorem also holds for any open subset of dimension d of a proper and smooth k -scheme.

Let X be an arbitrary reduced and separated k -scheme of dimension d . Using again (4.15) and the induction hypothesis, we may replace X by an arbitrarily small neighbourhood U of the generic points of its irreducible components. As the theorem holds for a scheme if and only if it holds for each of its connected components, we may shrink even more and assume that U is irreducible and affine. Let now $U \hookrightarrow X$ be an open immersion of U in a projective k -scheme X , such that U is dense in X . Using again the same argument, it suffices to prove the theorem for X .

Since X is projective, we can choose a closed immersion of X into a smooth formal W -scheme \mathbb{P} endowed with a Frobenius lifting F . Let N be an integer such that $N \geq 2d$. Then we can choose a proper hypercovering X_\bullet of X , a \mathbb{P} -augmented simplicial formal scheme \mathbb{P}_\bullet endowed with a Frobenius lifting F_\bullet , and a closed immersion of augmented simplicial schemes $X_\bullet \hookrightarrow \mathbb{P}_\bullet$ above $X \hookrightarrow \mathbb{P}$, so that the conditions of Lemma 5.1 are satisfied.

In particular, there is a dense open subset $U \subset X$ such that the restriction U_\bullet of X_\bullet above U is an étale hypercovering of U , and it suffices to prove the theorem for U . Let $Y = X \setminus U$. Since each \mathbb{P}_n is smooth over W , the complexes $(\mathcal{A}_{X_n, \mathbb{P}_n} \otimes \Omega_{\mathbb{P}_n}^\bullet \rightarrow \mathcal{A}_{Y_n, \mathbb{P}_n} \otimes \Omega_{\mathbb{P}_n}^\bullet)_t$ are defined. By functoriality, they define a complex of sheaves on X_\bullet [12, (5.1.6)], and we set

$$H_{\text{rig},c}^*(U_\bullet/K) := H^*(X_\bullet, (\mathcal{A}_{X_\bullet, \mathbb{P}_\bullet} \otimes \Omega_{\mathbb{P}_\bullet}^\bullet \rightarrow \mathcal{A}_{Y_\bullet, \mathbb{P}_\bullet} \otimes \Omega_{\mathbb{P}_\bullet}^\bullet)_t).$$

As usual, there is a biregular spectral sequence relating the cohomology of the individual complexes on the X_n 's to the cohomology of the global complex on X_\bullet , which can be written here

$$E_1^{i,j} = H_{\text{rig},c}^j(U_i/K) \Rightarrow H_{\text{rig},c}^{i+j}(U_\bullet/K).$$

By functoriality, this spectral sequence is endowed with a Frobenius automorphism. Moreover, the $E_1^{i,j}$ terms are finite dimensional K -vector spaces, so the spaces $H_{\text{rig},c}^{i+j}(U_\bullet/K)$ are finite dimensional too. Since the subspace of slope < 1 is an exact functor on the category of

finite dimensional K -vector spaces endowed with a σ -semilinear automorphism, we obtain a biregular spectral sequence

$$(5.3) \quad E_1^{i,j} = H_{\text{rig},c}^j(U_i/K)^{<1} \Rightarrow H_{\text{rig},c}^{i+j}(U_\bullet/K)^{<1}.$$

Let $\mathcal{I} \subset \mathcal{O}_X$ be a coherent ideal such that $V(\mathcal{I}) = Y$, and set $\mathcal{I}_n = \mathcal{I}\mathcal{O}_{X_n}$, $Y_n = V(\mathcal{I}_n) \subset X_n$. The sheaves $W\mathcal{O}_{X_n}$, $W\mathcal{O}_{Y_n}$ and $W\mathcal{I}_n$ define sheaves $W\mathcal{O}_{X_\bullet}$, $W\mathcal{O}_{Y_\bullet}$ and $W\mathcal{I}_\bullet$ on X_\bullet , with an exact sequence

$$0 \rightarrow W\mathcal{I}_\bullet \rightarrow W\mathcal{O}_{X_\bullet} \rightarrow W\mathcal{O}_{Y_\bullet} \rightarrow 0.$$

We define

$$H_c^*(U_\bullet, W\mathcal{O}_{U_\bullet,K}) = H^*(X_\bullet, W\mathcal{I}_\bullet,K),$$

and we obtain a biregular spectral sequence

$$(5.4) \quad E_1^{i,j} = H_c^j(U_i, W\mathcal{O}_{U_i,K}) \Rightarrow H_c^{i+j}(U_\bullet, W\mathcal{O}_{U_\bullet,K}).$$

Because F is an endomorphism of the simplicial scheme \mathbb{P}_\bullet , the homomorphisms (4.4) define morphisms of sheaves of rings $\mathcal{A}_{X_\bullet,\mathbb{P}_\bullet} \rightarrow W\mathcal{O}_{X_\bullet,K}$ and $\mathcal{A}_{Y_\bullet,\mathbb{P}_\bullet} \rightarrow W\mathcal{O}_{Y_\bullet,K}$ on X_\bullet , from which we derive a morphism of complexes

$$(\mathcal{A}_{X_\bullet,\mathbb{P}_\bullet} \otimes \Omega_{\mathbb{P}_\bullet}^\bullet \rightarrow \mathcal{A}_{Y_\bullet,\mathbb{P}_\bullet} \otimes \Omega_{\mathbb{P}_\bullet}^\bullet)_t \rightarrow (W\mathcal{O}_{X_\bullet,K} \rightarrow W\mathcal{O}_{Y_\bullet,K}).$$

Taking cohomology, we obtain homomorphisms

$$b_{U_\bullet}^q : H_{\text{rig},c}^q(U_\bullet/K)^{<1} \rightarrow H_c^q(U_\bullet, W\mathcal{O}_{U_\bullet,K}),$$

and a morphism of spectral sequences from (5.3) to (5.4). On the $E_1^{i,j}$ terms, this morphism is given by the homomorphisms $b_{U_i}^j$.

Since the truncation $\text{sk}_N^X(X_\bullet)$ is a N -truncated resolution of X , X_n is a projective and smooth k -scheme for all $n \leq N$. Then the theorem holds for the open subset $U_n \subset X_n$, which has dimension d since it is étale over U . Therefore, the homomorphisms $b_{U_i}^j$ between the $E_1^{i,j}$ terms of (5.3) and (5.4) are isomorphisms for $i \leq N$. It follows that $b_{U_\bullet}^q$ is an isomorphism for $q \leq N$.

As the augmentation $\mathbb{P}_\bullet \rightarrow \mathbb{P}$ is compatible with the Frobenius liftings, we obtain for all q a commutative square

$$(5.5) \quad \begin{array}{ccc} H_{\text{rig},c}^q(U_\bullet/K)^{<1} & \xrightarrow{b_{U_\bullet}^q} & H_c^q(U_\bullet, W\mathcal{O}_{U_\bullet,K}) \\ \uparrow & & \uparrow \\ H_{\text{rig},c}^q(U/K)^{<1} & \xrightarrow{b_U^q} & H_c^q(U, W\mathcal{O}_{U,K}). \end{array}$$

The augmented simplicial schemes $X_\bullet \rightarrow X$ and $Y_\bullet \rightarrow Y$ are proper hypercoverings, and the condition (iv) of 5.1 implies that $(X_\bullet, X_\bullet, \mathbb{P}_\bullet)$ and $(Y_\bullet, Y_\bullet, \mathbb{P}_\bullet)$ are Pr-Pr-Sm-hypercoverings of (X, X, \mathbb{P}) and (Y, Y, \mathbb{P})

in the sense of [28, 5.1.5]. Therefore, it follows from Tsuzuki's proper descent theorem ([27, 2.1.3], [28, 5.3.1]) that the left vertical arrow is an isomorphism. On the other hand, the right vertical arrow is an isomorphism thanks to Theorem 3.2. Since the source and target of b_U^q are 0 for $q > 2d$, and b_U^q is an isomorphism for $q \leq 2d$, this completes the proof of Theorem 1.1. \square

6. APPLICATIONS AND EXAMPLES

In this section, we assume that k is a finite field with $q = p^a$ elements, and we give applications of Theorem 1.1 to congruences mod q for the number of rational points of some algebraic varieties.

If X is a separated k -scheme of finite type, we set for all i

$$P_i(X, t) = \det(1 - t\phi|H_{\text{rig},c}^i(X/K)),$$

where $\phi = F^a$ is the k -linear Frobenius endomorphism of X . We recall that, thanks to the Lefschetz trace formula [16], the zeta function of X can be expressed as

$$(6.1) \quad \zeta(X, t) = \prod_i P_i(X, t)^{(-1)^{i+1}}.$$

6.1. We normalize the valuation on K by setting $v_q(q) = 1$. If $P(t) = \sum_i a_i t^i \in K[t]$, its Newton polygon is the graph of the greatest convex function s on $[0, \deg(P)]$ such that $s(i) \leq v_q(a_i)$ for all i . If P is irreducible, then its Newton polygon is a segment. For any polynomial $P(t) \in 1 + tK[t]$, and any $\lambda \in \mathbb{Q}$, we denote by $P^\lambda(t)$ the product of the irreducible factors of P whose Newton polygon is a segment of slope λ , normalised by $P^\lambda(0) = 1$. For any $\rho \in \mathbb{R}$, we set $P^{<\rho}(t) = \prod_{\lambda < \rho} P^\lambda(t)$.

This definition extends by multiplicativity to rational fractions $R(t) \in K(t)$ such that $R(0) = 1$. In particular, it can be applied to $\zeta(X, t) \in \mathbb{Q}(t) \subset K(t)$, for any separated k -scheme of finite type X , and this defines the *slope $< \rho$ factor* $\zeta^{<\rho}(X, t)$ of $\zeta(X, t)$. By Manin's theorem relating over finite fields the slopes with the eigenvalues of ϕ , we obtain

$$\det(1 - t\phi|H_{\text{rig},c}^i(X/K)^{<\rho}) = \det(1 - t\phi|H_{\text{rig},c}^i(X/K))^{<\rho}.$$

On the other hand, we define

$$P_i^W(X, t) = \det(1 - t\phi|H_c^i(X, W\mathcal{O}_{X,K})),$$

$$\zeta^W(X, t) = \prod_i P_i^W(X, t)^{(-1)^{i+1}}.$$

Then Theorem 1.1 implies Corollary 1.3:

$$(6.2) \quad \forall i, \quad P_i^{<1}(X, t) = P_i^W(X, t), \\ \zeta^{<1}(X, t) = \zeta^W(X, t).$$

We observe that these polynomials actually have coefficients in \mathbb{Z}_p :

Proposition 6.2. *For any separated k -scheme of finite type X , and any $\lambda \in \mathbb{Q}$, the polynomials $P_i^\lambda(X, t)$ and $P_i^W(X, t)$ belong to $\mathbb{Z}_p[t]$.*

Proof. By construction, the polynomials $P_i(X, t)$ belong to $K[t]$. Let $\phi' : X' \rightarrow X'$ be the pull-back of ϕ by the absolute Frobenius endomorphism of $\text{Spec}(k)$. By base change, we obtain

$$\det(1 - t\phi'|H_{\text{rig},c}^i(X'/K)) = \sigma^*(P_i(X, T)).$$

On the other hand, the relative Frobenius $F_{X/k} : X \rightarrow X'$ commutes with ϕ and ϕ' . As it induces an isomorphism $H_{\text{rig},c}^i(X'/K) \xrightarrow{\sim} H_{\text{rig},c}^i(X/K)$, it follows that

$$\det(1 - t\phi'|H_{\text{rig},c}^i(X'/K)) = P_i(X, T).$$

Thus $P_i(X, t)$ is invariant under σ , hence belongs to $\mathbb{Q}_p[t]$. Moreover, since the Newton polygon of a polynomial does not change by enlarging the field containing its coefficients, the decomposition $P_i(X, t) = \prod_\lambda P_i^\lambda(X, t)$ is defined in $\mathbb{Q}_p[t]$.

Finally, the slopes of the Frobenius action on $H_{\text{rig},c}^i(X/K)$ are non-negative [9, 3.1.2], and this is equivalent by Manin's theorem to the fact that the inverses of the roots of $P_i(X, t)$ are p -adic integers. Therefore, each $P_i^\lambda(X, t)$ belongs to $\mathbb{Z}_p[t]$.

By (6.2), the statement for $P_i^W(X, t)$ follows (it can also be proved directly by the same argument). \square

We recall the following well-known result [1], which links congruences modulo some power of q to the triviality of the corresponding slope factor:

Proposition 6.3. *Let $(N_r)_{r \geq 1}$ be a sequence of integers such that*

$$\Phi(t) = \exp\left(\sum_{r \geq 1} N_r \frac{t^r}{r}\right) \in 1 + \mathbb{Z}[[t]],$$

and assume that $\Phi(t)$ is a rational function in $\mathbb{Q}(t)$. Then $\Phi(t)$ can be written

$$\Phi(t) = \frac{\prod_i (1 - \alpha_i t)}{\prod_j (1 - \beta_j t)},$$

where $\alpha_i, \beta_j \in \overline{\mathbb{Z}}$ are such that $\alpha_i \neq \beta_j$ for all (i, j) , and, for any integer $\kappa \geq 1$, the following conditions are equivalent:

- (i) For all $r \geq 1$, $N_r \equiv 0 \pmod{q^{\kappa r}}$.
- (ii) For all i and all j , α_i and β_j are divisible by q^κ in $\overline{\mathbb{Z}}$.
- (iii) For all embeddings ι of $\overline{\mathbb{Q}}$ in $\overline{\mathbb{Q}}_p$, all i and all j , $v_q(\iota(\alpha_i)) \geq \kappa$, $v_q(\iota(\beta_j)) \geq \kappa$.
- (iv) $\Phi^{<\kappa}(t) = 1$.

As explained in the introduction, our first application of Theorem 1.1 over finite fields will be a proof of a consequence of Serre's conjecture on theta divisors.

Proposition 6.4. *Let X be a smooth projective variety of pure dimension n over a finite field. Let $D \subset X$ be an ample divisor, of complement $U = X \setminus D$, and assume that the following conditions hold:*

- a) *The injection $H^0(X, \omega_X) \rightarrow H^0(X, \omega_X(D))$ is an isomorphism;*
- b) *For all $i \geq 1$, the canonical homomorphism*

$$H^n(X, \mathcal{O}_X) \xrightarrow{V^i} H^n(X, W_{i+1}\mathcal{O}_X)$$

is injective.

Then $H_{\text{rig},c}^n(U/K)^{<1} \rightarrow H_{\text{rig}}^n(X/K)^{<1}$ is an isomorphism, and

$$(6.3) \quad \zeta^{<1}(X, t) = \zeta^{<1}(D, t) \cdot P_n^{<1}(X, t)^{(-1)^{n+1}}.$$

Proof. Let $\mathcal{I} \cong \mathcal{O}_X(-D)$ be the ideal of D in X . Condition a) is equivalent by Serre duality to the condition

$$H^n(X, \mathcal{I}) \xrightarrow{\sim} H^n(X, \mathcal{O}_X).$$

Thanks to condition b), the rows of the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^n(X, \mathcal{I}) & \longrightarrow & H^n(X, W_{i+1}\mathcal{I}) & \longrightarrow & H^n(X, W_i\mathcal{I}) \longrightarrow 0 \\ & & \downarrow \wr & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^n(X, \mathcal{O}_X) & \longrightarrow & H^n(X, W_{i+1}\mathcal{O}_X) & \longrightarrow & H^n(X, W_i\mathcal{O}_X) \longrightarrow 0 \end{array}$$

are exact, hence the homomorphism $H^n(X, W_i\mathcal{I}) \rightarrow H^n(X, W_i\mathcal{O}_X)$ is an isomorphism for all i . Taking inverse limits and tensoring with K , we obtain the isomorphism

$$(6.4) \quad H_c^n(U, W\mathcal{O}_{U,K}) \xrightarrow{\sim} H^n(X, W\mathcal{O}_{X,K}).$$

Thus Theorem 1.1 implies that $H_{\text{rig},c}^n(U/K)^{<1} \rightarrow H_{\text{rig}}^n(X/K)^{<1}$ is also an isomorphism.

The multiplicativity of the zeta function shows that it suffices to prove that

$$\zeta^{<1}(U, t) = P_n^{<1}(X, t)^{(-1)^{n+1}}.$$

By (6.2), we have

$$P_i^{<1}(U, t) = P_i^W(U, t)$$

for all i . By (6.4), we have $P_n^W(U, t) = P_n^W(X, t) = P_n^{<1}(X, t)$. On the other hand, U is affine and smooth of pure dimension n , hence $H_c^i(U, W\mathcal{O}_{U, K}) = 0$ for $i \neq n$ by Corollary 1.2. The proposition follows. \square

6.5. Proof of Theorem 1.4. Applying Proposition 6.3, it suffices to prove that, if Θ and Θ' are two theta divisors in an abelian variety A , then $\zeta^{<1}(\Theta, t) = \zeta^{<1}(\Theta', t)$. This will follow from (6.3) if we check that conditions a) and b) of Proposition 6.4 hold for a theta divisor in an abelian variety. Since A is an abelian variety, the dualizing bundle ω_A is trivial, and, for a theta divisor Θ , the homomorphism $H^0(A, \mathcal{O}_A) \rightarrow H^0(A, \mathcal{O}_A(\Theta))$ is an isomorphism [23, III 16]. Thus condition a) is satisfied. As for condition b), the injectivity of the homomorphisms $H^n(A, \mathcal{O}_A) \rightarrow H^n(A, W_{i+1}\mathcal{O}_A)$ follows from Serre's theorem on the vanishing of Bockstein operations for abelian varieties in characteristic p ([24, §1, 3] and [25, Théorème 2]). \square

Remarks 6.6. (i) In fact, Serre's conjecture is phrased more motivically: the difference of the motives of Θ and Θ' should be divisible in a suitable sense by the Lefschetz motive. What we show in Theorem 1.4 is a finite field implication of the motivic assertion. The motivic statement is clearly stronger than this implication. On the other hand, it is hard to approach directly as it deals with non-effective motives.

(ii) The property of the Theorem is very special for theta divisors. It is of course not true in general that two effective divisors D, D' with $h^0(\mathcal{O}(D)) = h^0(\mathcal{O}(D'))$ carry the same number of points modulo q : take for example $D = 2\{0\}, D' = \{0\} + \{\infty\}$ on \mathbb{P}^1 .

(iii) Let us also remark that over the complex numbers, when Θ is irreducible, one knows that its singularities are rational [13, Theorem 3.3]. A precise analogy of this over a finite field isn't quite clear, as the notion of rational singularities itself requires resolution of singularities. But it should be related to the assertion of Corollary 1.3.

According to Grothendieck-Deligne's philosophy of motives, Hodge type over the field of complex numbers behaves the same way as congruences for the number of rational points. In view of the previous remarks, it is worth pointing out that Theorem 1.4 has the following Hodge theoretic analogue (where $H^*(X)$ denotes the classical complex cohomology of an algebraic scheme X over \mathbb{C}):

Proposition 6.7. *Let A be an abelian variety of dimension n over \mathbb{C} , and $\Theta \subset A$ a theta divisor. Let F be the Deligne's Hodge filtration on $H^i(A)$ and $H^i(\Theta)$. Then the restriction map $\mathrm{gr}_F^0 H^i(A) \rightarrow \mathrm{gr}_F^0 H^i(\Theta)$ is*

an isomorphism for $i \neq n$. In particular, if Θ' is another theta divisor, $\dim \operatorname{gr}_F^0 H^i(\Theta) = \dim \operatorname{gr}_F^0 H^i(\Theta')$ for all i .

Proof. This is a simple example of application of [14, Proposition 1.2]. Since Θ is ample, we know by the weak Lefschetz theorem that the restriction map $H^i(A) \rightarrow H^i(\Theta)$ is an isomorphism for $i \leq n - 2$, and is injective for $i = n - 1$. Also $\operatorname{gr}_F^0 H^i(X) = 0$ for $i > \dim(X)$ for any separated \mathbb{C} -scheme of finite type X [12, Th. 8.2.4]. Thus all we have to prove is

$$(6.5) \quad \operatorname{gr}_F^0 H_c^n(U) \xrightarrow{\sim} \operatorname{gr}_F^0 H^n(A),$$

with $U = A \setminus \Theta$. Let $\sigma : A' \rightarrow A$ be a birational morphism such that $\sigma|_U$ is an isomorphism, A' is smooth, and $D = \sigma^{-1}(\Theta)_{\text{red}}$ is a normal crossings divisor. Then one has $\operatorname{gr}_F^0 H_c^n(U) = H^n(A', \mathcal{O}_{A'}(-D))$, and $\sigma^* : \operatorname{gr}_F^0 H^n(A) = H^n(A, \mathcal{O}_A) \rightarrow \operatorname{gr}_F^0 H^n(A') = H^n(A', \mathcal{O}_{A'})$ is an isomorphism. On the other hand, the composed morphism

$$(6.6) \quad H^n(A, \mathcal{O}_A(-\Theta)) \xrightarrow{\sigma^*} H^n(A', \mathcal{O}_{A'}(-D)) \xrightarrow{\iota} H^n(A', \mathcal{O}_{A'}) \xleftarrow{\sim} H^n(A, \mathcal{O}_A)$$

is an isomorphism since Θ is a theta divisor, while σ^*, ι are surjective for dimension reasons. Thus all maps in (6.6) are isomorphisms, which in particular proves the proposition. \square

We now consider the case of intersections of hypersurfaces of small degrees.

Proposition 6.8. *Let k be a perfect field of characteristic p , let $K = \operatorname{Frac}(W(k))$, and let $D_1, \dots, D_r \subset \mathbb{P}_k^n$ be hypersurfaces of degrees d_1, \dots, d_r . Assume that $\sum_j d_j \leq n$. Then, if $Z = D_1 \cap \dots \cap D_r$,*

$$(6.7) \quad H^0(Z, W\mathcal{O}_{Z,K}) = K, \quad H^i(Z, W\mathcal{O}_{Z,K}) = 0 \quad \text{for } i \geq 1.$$

Proof. We proceed by induction on r . Let $r = 1$, $D = D_1 = Z$, $d = d_1$. Then the condition $d \leq n$ implies that the homomorphisms $H^i(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}) \rightarrow H^i(D, \mathcal{O}_D)$ are isomorphisms for all i . It follows that, for any $m \geq 1$,

$$H^0(D, W_m \mathcal{O}_D) = W_m(k), \quad H^i(D, W_m \mathcal{O}_D) = 0 \quad \text{for } i \geq 1,$$

which implies (6.7).

For arbitrary r , let $D = D_1 \cup \dots \cup D_r$, which is a hypersurface of degree $d = \sum_j d_j \leq n$. For each sequence $1 \leq i_0 < \dots < i_s \leq r$, let $Z_{i_0, \dots, i_s} = D_{i_0} \cap \dots \cap D_{i_s}$. By Corollary 2.3, we obtain an exact sequence

$$0 \rightarrow W\mathcal{O}_{D,K} \rightarrow \prod_{i=1}^r W\mathcal{O}_{Z_i,K} \rightarrow \dots \rightarrow \prod_{i=1}^r W\mathcal{O}_{Z_{1, \dots, \hat{i}, \dots, r}, K} \rightarrow W\mathcal{O}_{Z,K} \rightarrow 0.$$

Applying the previous result to D , and the induction hypothesis to all Z_{i_0, \dots, i_j} for $j \leq r - 2$, we can view this exact sequence as providing a $\Gamma(\mathbb{P}^n, -)$ -acyclic left resolution of $W\mathcal{O}_{Z,K}$. Taking sections and observing that the complex

$$0 \rightarrow K \rightarrow \prod_{i=1}^r K \rightarrow \cdots \rightarrow \prod_{i=1}^r K \rightarrow 0$$

is acyclic, except in degree 0 where its cohomology is equal to K , we obtain (6.7). \square

Proof of Corollary 1.5. Combining Proposition 6.8 with Corollary 1.3, we obtain

$$\zeta^{<1}(Z, t) = \frac{1}{1-t},$$

which, by Proposition 6.3, is equivalent to congruence (1.7) for all finite extensions of \mathbb{F}_q . \square

We now discuss some cases where, given a morphism $f : X \rightarrow Y$ between two varieties over a finite field, Theorem 1.1 provides congruences between the numbers of rational points on X and Y .

Proposition 6.9. (i) *Let $f : X \rightarrow Y$ be a proper morphism between two separated \mathbb{F}_q -schemes of finite type. If the induced homomorphisms $f^* : H_c^i(Y, W\mathcal{O}_{Y,K}) \rightarrow H_c^i(X, W\mathcal{O}_{X,K})$ are isomorphisms for all $i \geq 0$, then*

$$(6.8) \quad |X(\mathbb{F}_q)| \equiv |Y(\mathbb{F}_q)| \pmod{q}.$$

(ii) *Let X be a proper scheme over \mathbb{F}_q , and G a finite group acting on X so that each orbit is contained in an affine open subset. If the action of G on $H^i(X, W\mathcal{O}_{X,K})$ is trivial for all i , then*

$$(6.9) \quad |X(\mathbb{F}_q)| \equiv |(X/G)(\mathbb{F}_q)| \pmod{q}.$$

Proof. Assertion (i) is an immediate consequence of 1.3 and 6.3, and implies assertion (ii) thanks to the following lemma:

Lemma 6.10. *Let X be a proper scheme over \mathbb{F}_q , G a finite group acting on X so that each orbit is contained in an affine open subset, $f : X \rightarrow Y := X/G$ the quotient map. Then f^* induces canonical isomorphisms*

$$(6.10) \quad H^i(Y, W\mathcal{O}_{Y,K}) \xrightarrow{\sim} H^i(X, W\mathcal{O}_{X,K})^G$$

for all i .

The canonical morphism $f : X \rightarrow Y$ is finite, and $\mathcal{O}_Y \xrightarrow{\sim} f_*(\mathcal{O}_X)^G$. By induction on m , $R^i f_*(W_m \mathcal{O}_X) = 0$ for all $i \geq 1$ and all $m \geq 1$. Moreover, the morphisms

$$W_m \mathcal{O}_Y \rightarrow W_m(f_*(\mathcal{O}_X)^G) \rightarrow W_m(f_*(\mathcal{O}_X))^G \cong f_*(W_m \mathcal{O}_X)^G$$

are isomorphisms for all $m \geq 1$. Since taking invariants under G commutes with inverse limits and with tensorisation by \mathbb{Q} , they provide isomorphisms

$$W\mathcal{O}_{Y,K} \xrightarrow{\sim} (f_*(W\mathcal{O}_X)^G)_K \xrightarrow{\sim} f_*(W\mathcal{O}_{X,K})^G.$$

As $\text{char}(K) = 0$, taking invariants under G commutes with cohomology for $K[G]$ -modules, and we can write

$$H^i(Y, W\mathcal{O}_{Y,K}) \xrightarrow{\sim} H^i(Y, f_*(W\mathcal{O}_{X,K})^G) \xrightarrow{\sim} H^i(Y, f_*(W\mathcal{O}_{X,K}))^G.$$

On the other hand, the inverse system $(f_*(W_m \mathcal{O}_X))_{m \geq 1}$ is \varprojlim -acyclic, and f_* commutes with tensorisation with K . Therefore the morphism

$$f_*(W\mathcal{O}_{X,K}) \rightarrow \mathbb{R}f_*(W\mathcal{O}_{X,K})$$

is an isomorphism, and we obtain

$$H^i(Y, f_*(W\mathcal{O}_{X,K}))^G \xrightarrow{\sim} H^i(X, W\mathcal{O}_{X,K})^G,$$

which proves the lemma. \square

6.11. In the most favorable cases, the assumptions of the previous proposition can be checked in characteristic p . This is the case under the assumptions of Corollary 1.6, the proof of which follows:

Proof of Corollary 1.6. Let $f : X \rightarrow Y$ be a morphism between two proper \mathbb{F}_q -schemes such that the induced homomorphisms $f^* : H^i(Y, \mathcal{O}_Y) \rightarrow H^i(X, \mathcal{O}_X)$ are isomorphisms for all $i \geq 0$. By induction on m , it follows that $H^i(Y, W_m \mathcal{O}_Y) \rightarrow H^i(X, W_m \mathcal{O}_X)$ is an isomorphism for all $i \geq 0$ and all $m \geq 1$. As X and Y are proper, we obtain that

$$H^i(Y, W\mathcal{O}_{Y,K}) \xrightarrow{\sim} H^i(X, W\mathcal{O}_{X,K})$$

for all $i \geq 0$. Then congruence (1.8) again follows from 1.3 and 6.3. \square

Corollary 6.12. *Let X be a proper scheme over \mathbb{F}_q , and G a finite group acting on X so that each orbit is contained in an affine open subset of X . If $|G|$ is prime to p , and if the action of G on $H^i(X, \mathcal{O}_X)$ is trivial for all i , then*

$$(6.11) \quad |X(\mathbb{F}_q)| \equiv |(X/G)(\mathbb{F}_q)| \pmod{q}.$$

Proof. When $|G|$ is prime to p , taking invariants under G is an exact functor on $k[G]$ -modules, and we obtain isomorphisms

$$\begin{aligned} H^i(Y, \mathcal{O}_Y) &\xrightarrow{\sim} H^i(Y, (\mathbb{R}f_*\mathcal{O}_X)^G) \xrightarrow{\sim} H^i(Y, (\mathbb{R}f_*\mathcal{O}_X))^G \\ &\xrightarrow{\sim} H^i(X, \mathcal{O}_X)^G = H^i(X, \mathcal{O}_X). \end{aligned}$$

Hence we can apply Corollary 1.6. \square

Remarks 6.13. Without assumption on $|G|$, Fu and Wan have proved [30, Theorem 0.1] that the congruence (6.9) holds under the following hypotheses:

- a) X is the reduction of a projective and smooth W -scheme X' with p -torsion free Hodge cohomology;
- b) The action of G on X is the reduction of an action on X' such that the induced action on $H^i(X', \mathcal{O}_{X'})$ is trivial for all i .

We do not know whether, under these hypotheses, the action of G is trivial on the spaces $H^i(X, W\mathcal{O}_{X,K})$. However, the following example seems to indicate that, when p divides $|G|$, a mod p assumption on the action of G as in 6.12 does not suffice to provide congruences such as (6.9).

Let $p = 2$, let E_1 be an elliptic curve over \mathbb{F}_q with an \mathbb{F}_q -rational point t of order 2, and let E_2 be another elliptic curve over \mathbb{F}_q . We define $X = E_1 \times E_2$, and we let the group $G = \mathbb{Z}/2\mathbb{Z}$ act on X via $(x, y) \mapsto (x+t, -y)$, so that $Y = X/G$ is the classical Igusa surface [19], which is smooth over \mathbb{F}_q . The action of G on $H^i(X, \mathcal{O}_X)$ is trivial for all i , but, thanks to the Künneth formula in crystalline cohomology, one checks easily that $H^2(X, W\mathcal{O}_{X,K}) \cong H^2(X/K)^{<1} \neq 0$, while $H^2(Y, W\mathcal{O}_{Y,K}) \cong (H^2(X/K)^G)^{<1} = 0$. Therefore $\zeta^{<1}(Y, t) \neq \zeta^{<1}(X, t)$, and a congruence such as (6.9) cannot hold for all powers of q .

Note also that, for any p , taking the quotient of an abelian variety by the subgroup generated by a rational point of order p provides an example where the assumptions of 6.9 (ii) are satisfied, while those of 1.6 fail to be true.

REFERENCES

- [1] Ax, J.: Zeroes of polynomials over finite fields, *Am. J. of Mathematics* **86** (1964), 255–261.
- [2] Berthelot, P.: Géométrie rigide et cohomologie des variétés algébriques de caractéristique p , Journées d’Analyse p -adique (Luminy, 1982), in *Introduction aux cohomologies p -adiques*, Mémoires Soc. Math. France **23** (1986), 7–32.
- [3] Berthelot, P.: Cohomologie rigide et cohomologie rigide à supports propres, première partie, Prépublication IRMAR 96-03, Université de Rennes (1996).
- [4] Berthelot, P.: Finitude et pureté cohomologique en cohomologie rigide, avec un appendice par A. J. de Jong, *Invent. math.* **128** (1997), 329–377.
- [5] Berthelot, P.: Dualité de Poincaré et formule de Künneth en cohomologie rigide, *Comptes-Rendus Acad. Sc. Paris* **325** (1997), 493–498.
- [6] Bloch, S.: Algebraic K -theory and crystalline cohomology, *Publ. math. IHÉS* **47** (1977), 187–268.
- [7] Bloch, S., Esnault, H., Levine, M.: Decomposition of the diagonal and eigenvalues of Frobenius for Fano hypersurfaces, *Am. J. of Mathematics* **127** (2005), 193–207.
- [8] Bloom, T., Herrera, M.: De Rham cohomology of an analytic space, *Invent. math.* **7** (1969), 275–296.
- [9] Chiarellotto, B., Le Stum, B.: Pentes en cohomologie rigide et F -isocristaux unipotents, *Manuscripta Math.* **100** (1999), 455–468.
- [10] Chiarellotto, B., Tsuzuki, N.: Cohomological descent of rigid cohomology for étale coverings, *Rend. Sem. Mat. Univ. Padova* **109** (2003), 63–215.
- [11] de Jong, J.: Smoothness, semi-stability and alterations, *Publ. math. IHÉS* **83** (1996), 51–93.
- [12] Deligne, P.: Théorie de Hodge III, *Publ. math. IHÉS* **44** (1974), 5–77.
- [13] Ein, L., Lazarsfeld, R.: Singularities of theta divisors and the birational geometry of irregular varieties, *J. Amer. Math. Soc.* **10** (1997), 1, 243–258.
- [14] Esnault, H.: Hodge type of subvarieties of \mathbb{P}^n of small degrees, *Math. Ann.* **288** (1990), 549–551.
- [15] Esnault, H.: Appendix to “Congruences for rational points on varieties defined over finite fields” by N. Fakhruddin and C. S. Rajan, appears in *Math. Ann.* (2005).
- [16] Étesse, J.-Y.; Le Stum, B.: Fonctions L associées aux F -isocristaux surconvergens. I, *Math. Ann.* **296** (1993), 557–576.
- [17] Grothendieck, A.: Éléments de géométrie algébrique, *Publ. math. IHÉS* **11** (1961), 5–167.
- [18] Hartshorne, R.: *Residues and Duality*, Lecture Notes in Mathematics **20** (1966), Springer-Verlag.
- [19] Igusa, J.-I.: On some problems in abstract algebraic geometry, *Proc. Nat. Acad. Sci. U. S. A.* **41** (1955), 964–967.
- [20] Illusie, L.: Complexe de de Rham-Witt et cohomologie cristalline, *Ann. Éc. Norm. Sup.* **4** **12** (1979), 501–661.
- [21] Katz, N.: On a theorem of Ax, *Am. J. of Mathematics* **93** (1971), 485–499.
- [22] Kiehl, R.: Theorem A und Theorem B in der nichtarchimedischen Funktionentheorie, *Invent. math.* **2** (1967), 256–273.

- [23] Mumford, D.: Abelian Varieties, Tata Institute of Fundamental Research (1970).
- [24] Serre, J.-P.: Sur la topologie des variétés algébriques en caractéristique p , in *Symposium internacional de topología algebraica* (International symposium on algebraic topology), 24–53, Universidad Nacional Autonoma de Mexico and UNESCO, Mexico City, 1958.
- [25] Serre, J.-P.: Quelques propriétés des variétés abéliennes en caractéristique p , *Amer. J. Math.* **80** (1958), 715–739.
- [26] Serre, J.-P.: *Corps locaux*, Publ. Inst. Mathématique Univ. Nancago **8** (1962), Hermann.
- [27] Tsuzuki, N.: Cohomological descent of rigid cohomology for proper coverings, *Invent. math.* **151** (2003), 101–133.
- [28] Tsuzuki, N.: Cohomological descent in rigid cohomology, in *Geometric aspects of Dwork theory*, edited by A. Adolphson et alii, vol.2, De Gruyter (2004), 931–981.
- [29] Wan, D.: Mirror symmetry for zeta functions, preprint (2004).
- [30] Fu, L., Wan, D.: Mirror congruences for rational points on Calabi-Yau varieties, preprint (2005).

IRMAR, UNIVERSITÉ DE RENNES 1, CAMPUS DE BEAULIEU, 35042 RENNES
CEDEX, FRANCE

E-mail address: pierre.berthelot@univ-rennes1.fr

DEPT. OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, IL 60637,
USA

E-mail address: bloch@math.uchicago.edu

MATHEMATIK, UNIVERSITÄT DUISBURG-ESSEN, FB6, MATHEMATIK, 45117
ESSEN, GERMANY

E-mail address: esnault@uni-essen.de