

Algebraic Geometry

Coniveau over p -adic fields and points over finite fields

Hélène Esnault¹

Universität Duisburg-Essen, Mathematik, 45117 Essen, Germany

Received 26 April 2007; accepted 6 May 2007

Available online 5 July 2007

Presented by Pierre Deligne

Abstract

If the ℓ -adic cohomology of a projective smooth variety, defined over a p -adic field K with finite residue field k , is supported in codimension ≥ 1 , then any model over the ring of integers of K has a k -rational point. **To cite this article:** H. Esnault, C. R. Acad. Sci. Paris, Ser. I 345 (2007).

© 2007 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Résumé

Coniveau sur un corps p -adique et points sur un corps fini. Si la cohomologie ℓ -adique d'une variété projective, lisse, définie sur un corps p -adique K à corps résiduel fini k , est supportée en codimension ≥ 1 , alors tout modèle sur l'anneau des entiers de K a un point rationnel. **Pour citer cet article :** H. Esnault, C. R. Acad. Sci. Paris, Ser. I 345 (2007).

© 2007 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Version française abrégée

Soit X une variété projective et absolument irréductible sur un corps local K . Rappelons qu'un modèle de X/K sur l'anneau de valuation R de K est un morphisme $\mathcal{X} \rightarrow \text{Spec } R$ projectif et plat, tel que $(\mathcal{X} \rightarrow \text{Spec } R) \otimes K = (X \rightarrow \text{Spec } K)$. Nous considérons la cohomologie ℓ -adique $H^i(\bar{X})$ à coefficients dans \mathbb{Q}_ℓ . Dire qu'elle est supportée en codimension 1 signifie que toute classe dans $H^i(\bar{X})$ a une restriction nulle dans $H^i(\bar{U})$, où $U \subset X$ est un ouvert non vide. Le but de cette Note est de prouver le théorème suivant :

Théorème. Soit X une variété projective lisse et absolument irréductible sur un corps local K de caractéristique 0 et à corps résiduel fini k . On suppose que la cohomologie ℓ -adique $H^i(\bar{X})$ est supportée en codimension ≥ 1 pour tout $i \geq 1$. Soit \mathcal{X}/R un modèle. Alors il existe un morphisme projectif surjectif $\sigma : \mathcal{Y} \rightarrow \mathcal{X}$ de R -schémas tel que $|\mathcal{Y}(k)| \equiv 1 \text{ modulo } |k|$.

On en déduit immédiatement le corollaire suivant :

Corollaire. Sous les hypothèses du théorème, tout modèle \mathcal{X}/R possède un point k -rationnel.

E-mail address: esnault@uni-due.de.

¹ Partially supported by the DFG Leibniz Preis.

Pour ce qui concerne l’existence du point k -rationnel, ceci affranchit [3, Theorem 1.1], (qui est vrai aussi si K est de caractéristique $p > 0$), de l’hypothèse de régularité sur le choix du modèle \mathcal{X} , qui était utilisée pour pouvoir appliquer le théorème de pureté de Gabber [4]. Pour ce faire, nous montrons que d’avoir des singularités quotient est suffisant, de même que pour l’étude de l’application de spécialisation. Nous appliquons alors la version plus précise du théorème de Jong ainsi qu’elle est exposée dans [1].

1. Introduction

Let X be a projective, absolutely irreducible variety defined over a local field K with finite residue field k . Recall that a *model* of X/K on the valuation ring R of K is a flat projective morphism $\mathcal{X} \rightarrow \text{Spec } R$ such that $(\mathcal{X} \rightarrow \text{Spec } R) \otimes K = (X \rightarrow \text{Spec } K)$. We consider ℓ -adic cohomology $H^i(\bar{X})$ with \mathbb{Q}_ℓ -coefficients. One defines the first coniveau level

$$N^1 H^i(\bar{X}) = \{ \alpha \in H^i(\bar{X}), \exists \text{ divisor } D \subset X \text{ s.t. } 0 = \alpha|_{X \setminus D} \in H^i(\overline{X \setminus D}) \}. \tag{1}$$

As $H^i(\bar{X})$ is a finite dimensional \mathbb{Q}_ℓ -vector space, one has by localization

$$\exists D \subset X \text{ s.t. } N^1 H^i(\bar{X}) = \text{Im}(H^i_D(\bar{X}) \rightarrow H^i(\bar{X})), \tag{2}$$

where $D \subset X$ is a divisor. One says that $H^i(\bar{X})$ is *supported in codimension 1* if $N^1 H^i(\bar{X}) = H^i(\bar{X})$. This definition is general, but has good properties only if X is irreducible and smooth or has only very mild singularities.

In [3, Theorem 1.1] it is shown that if X/K is smooth, projective, absolutely irreducible over a local field K with finite residue field k , and if ℓ -adic cohomology $H^i(\bar{X})$ is supported in codimension ≥ 1 for all $i \geq 1$, then any regular model \mathcal{X}/R of X/K has the property

$$|\mathcal{X}(k)| \equiv 1 \pmod{|k|}. \tag{3}$$

The purpose of this Note is to drop the regularity assumption if K has characteristic 0.

Theorem 1.1. *Let X be a smooth, projective, absolutely irreducible variety defined over a local field K of characteristic 0 with finite residue field k . Assume that the ℓ -adic cohomology $H^i(\bar{X})$ is supported in codimension ≥ 1 for all $i \geq 1$. Let \mathcal{X} be a model of X over the ring of integers R of K . Then there is a projective surjective morphism $\sigma : \mathcal{Y} \rightarrow \mathcal{X}$ of R -schemes such that*

$$|\mathcal{Y}(k)| \equiv 1 \pmod{|k|}.$$

As an immediate corollary, one obtains:

Corollary 1.2. *Under the assumptions of the theorem, every model \mathcal{X}/R has a k -rational point.*

The regularity of the model \mathcal{X} in the proof of [3, Theorem 1.1] (which is shown also when K has characteristic $p > 0$) was used to apply Gabber’s purity theorem [4]. We show that for the piece of regularity one needs, it is enough to have quotient singularities. Likewise, for the properties needed on the specialization map, quotient singularities are good enough. The more careful use of de Jong’s theorem as exposed in [1] allows one then to conclude.

2. Proof of Theorem 1.1

Let K be a local field of characteristic 0 with finite residue field k . Let $R \subset K$ be its valuation ring. Let $\mathcal{X} \rightarrow \text{Spec } R$ be an integral model of a projective variety $X \rightarrow \text{Spec } K$. We do not assume here that X is absolutely irreducible, nor do we assume that X/K is smooth. Then by [1, Corollary 5.15], there is a diagram

$$\begin{array}{ccccc}
 \mathcal{Z} & \xrightarrow{\pi} & \mathcal{Y} & \xrightarrow{\sigma} & \mathcal{X} \\
 & & \searrow & & \downarrow \\
 & & & & \text{Spec } R
 \end{array} \tag{4}$$

and a finite group G acting on \mathcal{Z} over \mathcal{Y} with the properties

- (i) $\mathcal{Z} \rightarrow \text{Spec } R$ and $\mathcal{Y} \rightarrow \text{Spec } R$ are flat,
- (ii) σ is projective, surjective, and birational,
- (iii) \mathcal{Y} is the quotient of \mathcal{Z} by G ,
- (iv) \mathcal{Z} is regular.

So $\mathcal{Y} \rightarrow \text{Spec } R$ is not quite a model of $\mathcal{X} \rightarrow \text{Spec } K$, but is close to it. We show in the sequel that σ in (4) does it in Theorem 1.1. Set

$$Y = \mathcal{Y} \otimes K, \quad Z = \mathcal{Z} \otimes K.$$

For an open $U \subset X$ let us set $Y_U = U \times_X Y$, $Z_U = U \times_X Z$.

Let us assume now that X/K is smooth. This implies that

$$\sigma^* : H^i(\bar{U}) \hookrightarrow H^i(\bar{Y}_U) \tag{5}$$

is injective. Moreover, the trace map from Y to X

$$H^i(\bar{Y}_U) \xrightarrow{(\text{trace})_{Y/X}} H^i(\bar{U}) \tag{6}$$

is a retraction of σ^* in (5). Let $i \geq 1$ and let $D \subset X$ be a divisor such that $H^i_D(\bar{X}) \rightarrow H^i(\bar{X})$ and such that $\sigma|_{X \setminus D} : Y \setminus \sigma^{-1}(D) \rightarrow X \setminus D$ is an isomorphism. Then (6) yields the commutative diagram

$$\begin{array}{ccc} H^i(\bar{Y}) & \longrightarrow & H^i(\overline{Y \setminus \sigma^{-1}(D)}) \\ (\text{trace})_{Y/X} \downarrow & & \downarrow = \\ H^i(\bar{X}) & \xrightarrow{0} & H^i(\overline{X \setminus D}) \end{array} \tag{7}$$

and we conclude

$$X/K \text{ smooth} \implies N^1 H^i(\bar{X}) = H^i(\bar{X}) \subset N^1 H^i(\bar{Y}) = H^i(\bar{Y}). \tag{8}$$

We endow all schemes considered (which are R -schemes) with the upper subscript u to indicate the base change $\otimes_R R^u$ or $\otimes_K K^u$, where $K^u \supset K$ is the maximal unramified extension, and $R^u \supset R$ is the normalization of R in K^u . Likewise, we write $\bar{}$ to indicate the base change $\otimes_R \bar{R}$, $\otimes_K \bar{K}$, $\otimes_k \bar{k}$, where $\bar{K} \supset K$, $\bar{k} \supset k$ are the algebraic closures and $\bar{R} \supset R$ is the normalization of R in \bar{K} . We consider as in [3, (2.1)] the F -equivariant exact sequence [2, 3.6(6)]

$$\dots \rightarrow H^i_B(\mathcal{Y}^u) \xrightarrow{\iota} H^i(\bar{B}) = H^i(\mathcal{Y}^u) \xrightarrow{sp^u} H^i(Y^u) \rightarrow \dots, \tag{9}$$

where $F \in \text{Gal}(\bar{k}/k)$ is the geometric Frobenius, and $B = \mathcal{Y} \otimes k$.

One has

Claim 2.1. *The eigenvalues of the geometric Frobenius $F \in \text{Gal}(\bar{k}/k)$ acting on $H^i(X^u)$ and on $H^i(Y^u)$ lie in $q \cdot \bar{\mathbb{Z}}$ for all $i \geq 1$.*

Proof. For $H^i(X^u)$, this is [3, Theorem 1.5(ii)]. One has $H^i(\bar{Y}) = H^i(\bar{Z})^G$, thus in particular, $\pi^* : H^i(\bar{Y}) \rightarrow H^i(\bar{Z})$ is injective. By (8) one has

$$H^i(\bar{Y}) \xrightarrow{\pi^* \text{ inj}} N^1 H^i(\bar{Z}). \tag{10}$$

Since K has characteristic 0, and \mathcal{Z} is regular by (iv), Z is smooth. Thus we can apply again [3, Theorem 1.5(ii)]. This finishes the proof. \square

Claim 2.2. *The eigenvalues of the geometric Frobenius $F \in \text{Gal}(\bar{k}/k)$ acting on $\iota(H^i_C(\mathcal{Y}^u)) \subset H^i(\bar{B})$ lie in $q \cdot \bar{\mathbb{Z}}$ for all $i \geq 1$.*

Proof. By (iii), one has $H^i_B(\mathcal{Y}^u) = H^i_C(\mathcal{Z}^u)^G \subset H^i_C(\mathcal{Z}^u)$, where $C = \pi^{-1}(B)$. Since by (iv), \mathcal{Z} is regular, we can apply [3, Theorem 1.4], which is a consequence of Gabber’s purity theorem [4], to conclude. \square

Proof of Theorem 1.1. Claims 2.1 and 2.2 together with (9) show that the eigenvalues of F acting on $H^i(\bar{B})$ lie in $q \cdot \bar{\mathbb{Z}}$ for all $i \geq 1$.

We apply the Lefschetz trace formula $|B(k)| = \text{Tr } F|H^*(\bar{B})$. As B is absolutely connected and defined over k , $F|H^0(\bar{B}) = \text{Identity}$. By the discussion, one has $|B(k)| \in \mathbb{N} \cap (1 + q \cdot \bar{\mathbb{Z}}) \subset 1 + q \cdot \mathbb{Z}$. \square

3. Remarks

Starting from Theorem 1.1, and Corollary 1.2, we may ask what happens if K has equal characteristic $p > 0$ and whether or not the congruence of the theorem is true on all models. We have no counter-examples for either question. As far as K is concerned, characteristic 0 is used in the proof of Claim 2.1: if K has characteristic $p > 0$, we only know that Z is regular, thus we cannot apply immediately [3, Theorem 1.5(ii)]. Going up to a strict semi-stable model does not help as for this, one has to ramify R and one loses regularity of \mathcal{Z} and Z . As far as the congruence is concerned, instead of going to a birational model \mathcal{Y} (or birational up to some inseparable extension in characteristic $p > 0$), one should go up to a hypercover built out of such \mathcal{Y} . In doing Deligne's construction of hypercovers with resolutions of singularities being replaced by de Jong's morphisms of the type σ in (4), one creates components which do not dominate \mathcal{X} , the cohomology of which is very hard to control. So one perhaps loses the coniveau property.

Acknowledgements

This Note relies on de Jong's fundamental alteration theorems. T. Saito suggested to us the use of them in the shape formulated in [1]. We thank him for this, and for many subsequent discussions on the subject. We exposed a weaker version of Theorem 1.1 at the conference in honor of S. Bloch in Toronto in March 2007. Discussions with him, A. Beilinson and L. Illusie contributed to simplify our original exposition.

References

- [1] A.J. de Jong, Families of curves and alterations, *Ann. Inst. Fourier* 47 (2) (1997) 599–621.
- [2] P. Deligne, La conjecture de Weil, II, *Publ. Math. IHES* 52 (1981) 137–252.
- [3] H. Esnault, Deligne's integrality theorem in unequal characteristic and rational points over finite fields, with an appendix with P. Deligne, *Ann. Math.* 164 (2006) 715–730.
- [4] K. Fujiwara, A proof of the absolute purity conjecture (after Gabber), in: *Algebraic Geometry 2000, Azumino*, in: *Advanced Studies in Pure Mathematics*, vol. 36, Mathematical Society of Japan, 2002, pp. 153–183.